



ELSEVIER

Journal of Arid Environments 64 (2006) 422–435

www.elsevier.com/locate/jnlabr/yjare

Journal of
Arid
Environments

Testudo graeca graeca feeding ecology in an arid and overgrazed zone in Morocco

E.H. El Mouden^a, T. Slimani^a, K. Ben Kaddour^a, F. Lagarde^{b,*},
A. Ouhammou^c, X. Bonnet^b

^aLaboratoire d'Ecologie Animale Terrestre, Faculté des Sciences, Semlalia,
Université Cadi Ayyad, B.P. 2390, 40000 Marrakech, Maroc

^bCentre d'Etudes Biologiques de Chizé-CNRS, 79360 Villiers en Bois, France

^cLaboratoire d'Ecologie Végétale, Département de Biologie, Faculté des Sciences-Semlalia,
Université Cadi Ayyad, B.P. 2390, 40000 Marrakech, Maroc

Received 30 December 2004; received in revised form 3 June 2005; accepted 11 June 2005
Available online 20 July 2005

Abstract

Terrestrial chelonians of arid regions are frequently faced with highly variable food quality and/or quantity and these problems could be aggravated in overgrazing areas. Then, it is crucial for species conservation to possess precise data on feeding ecology. In this paper, we provide the first quantitative data on *Testudo graeca graeca* feeding ecology, in the central Jbilets (Morocco). Diet composition was obtained by faecal analysis and compared to plant assemblages in the field. Interestingly, *T. g. graeca* seems rather a specialist herbivorous tortoise. The five main plant species found in faecal samples (*Leontodon saxatilis* (Asteraceae); *Malva parviflora* (Malvaceae); *Astragalus cruciatus*, *Medicago hispida* and *Lotus arenarius* (Fabaceae)) represent 70% of the identified material and are highly preferred whereas several other plant species, common in the field (e.g. *Eryngium ilicifolium* (Ombellifereae); *Emex spinosus* (Polygonaceae); *Spergula flaccida* Caryophyllaceae)) are actively avoided. Fabaceae made up 27% of the diet and may be important forbs in the diet of terrestrial tortoise owing to their high nutritional value. The diet composition of *T. g. graeca* suggests that diet overlap may occur between domestic ungulates and tortoises in overgrazed landscape and could

*Corresponding author. Tel.: +33 549 097 840, +33 549 096 111; fax: +33 549 096 526.
E-mail address: lagarde@cebc.cnrs.fr (F. Lagarde).

generate a competition context. However, *T. g. graeca* seems to focus its foraging effort under the spiny shrubs where the impact of overgrazing is strongly attenuated.

© 2005 Elsevier Ltd. All rights reserved.

Keywords: Diet selection; Faecal analysis; Herbivory; Spur thighed tortoise; Competition

1. Introduction

The Mediterranean Spur-thighed tortoise *Testudo graeca graeca* is the unique terrestrial tortoise in Morocco. Populations have declined substantially by habitat changes due to human activities (e.g. agriculture, overgrazing, deforestation) (Bayley and Highfield, 1996; Slimani et al., 2001). Various studies have already been conducted on population dynamics (Bayley and Highfield, 1996; El Mouden et al., 2001; Slimani et al., 2001), general ecology (Lambert, 1969, 1981, 1983), geographic variations (Highfield, 1990a, b) and impact of trade (Highfield, 1994; Bayley and Highfield, 1996). Unfortunately, food habits are still poorly known and limited to brief notes in the literature (Cobo and Andreu, 1988; Highfield, 1989; Bayley and Highfield, 1996; Andreu et al., 2000). Precise qualitative and quantitative information about tortoise feeding ecology are nevertheless fundamental to conservation purpose in overgrazing area.

Several methods have been developed to evaluate the composition of herbivore diets: direct observation of animal foraging behaviour, and indirect methods of dietary reconstitution (alkane technique, stomach and faecal analysis). The direct observation of animals, when possible, is the more accurate sampling method but is exposed to several potential problems. It could be very difficult to identify precisely the items consumed by an individual and the presence of the observer can change the behaviour of animals (Gordon, 1995). This method is classically time-consuming and the observer can only sample a small number of individuals at a time (Gordon, 1995). This problem is particularly cumbersome to study Testudines species due to their sporadic activity and infrequent foraging behaviour (Lagarde et al., 2002, 2003). Nevertheless, direct observations of free ranging tortoises can be relatively straightforward and fruitful for diet and diet selection analysis, especially for desert tortoises (Jennings, 1993; Lagarde et al., 2003). Unfortunately, tortoises often forage under dense vegetation cover, and are therefore very difficult to observe. Then, indirect methods of feeding analysis provide helpful surrogates.

The faecal analysis method is the more widely used technique for determining herbivore diets indirectly (Baumgartner and Martin, 1939; Hercus, 1960; Stewart, 1967; Sparks and Malechek, 1968; Chapuis, 1980; Varva and Holechek, 1980; Garcia-Gonzalez, 1984; Macdonald and Mushinsky, 1988; Milton, 1992; Mason et al., 1999; Loehr, 2002). This technique can provide useful information on qualitative and quantitative aspects of herbivore diets (Chapuis and Didillon, 1987) despite the potential bias due to differential digestibility of plant species (Stewart, 1970; Varva et al., 1978; McInnis et al., 1983).

Then, the aim of our study is to give preliminary results about feeding ecology and diet diversity of *T. g. graeca* in an arid and overgrazed area in Morocco, using the faecal analysis technique.

2. Materials and methods

2.1. Study area

The study site is a 33 ha area located in the central Jbilet mountains about 25 km north of Marrakech, in the western of Morocco (31°37'N, 8°02'W, altitude: 580 m above sea level). The study site is located in an arid region, with mean annual rainfall of about 240 mm and most precipitation occurring between September and February (El Mouden et al., 1999; Znari et al., 2002). Average air temperature in the hottest month (July) can reach 39 °C and the minimal annual temperature is normally above 0 °C (Emberger, 1933; Le Houérou, 1989). Vegetation consists mainly in spreaded bushes of Jujube (*Ziziphus lotus*), with some widely separated Acacia (*Acacia gummifera*) (typical arid vegetation) and Retams (*Retama monosperma*). Most of habitat is open, hard bare ground with stony soils on the flats and low hillsides that surround small sandy, pebbly or stony wadis (riverbeds). Seasonal over-grazing by domestic livestock (sheep and goat) strongly affects the vegetation structure. A simple visual inspection of study area reveal two major and different plant assemblages: a relatively rich and diversified plant community under Jujube, protected against over-grazing by the spiny structure of this shrub, and a lesser diversified plant community out of the Jujube, with a low percent of plant cover.

2.2. Vegetal cover

The study was performed in spring 2002 (May–June), during the period of maximum activity for tortoises. All plant species represent in the study area were collected and identified at the species level, with the help of previous botanical studies (Nègre, 1961, 1962; Quezel and Santa, 1962, 1963).

To describe the vegetation structure, the percentage of plant cover for each species was measured in 43 quadrats (1 m²) placed at 100 m from each other along six parallel lines established in the study area. Nine quadrats were sampled under jujube shrubs, 34 quadrats were set up out of the jujube shrubs, broadly reflecting the actual proportion occupied by each major types of habitat in the study site.

2.3. Faecal analysis

Captured animals were considered as juveniles when midline carapace length was less than 100 mm (Slimani et al., 2001), the minimum size observed at maturity. Adult were sexed using classical dimorphic characters for *Testudo* (Andreu et al., 2000; Bonnet et al., 2001; El Mouden et al., 2001). The faecal samples were obtained during the animal's manipulation and collected with a metallic pincer disinfected

with alcohol in order to avoid diseases and parasites transmission between individuals.

To allow species recognition of faecal plant fragments, a reference collection of upper, lower leaves and caudal epidermis were realized. Epidermises were obtained by mechanical separation on fresh plant material with a scalpel under a microscope. Then, each fragment were placed in potassium chloride for 10 min, rinsed in distilled water, mounted on microscope slides and photographed under an optic microscope ($320\times$).

A total of 44 faeces were collected during the study period (seven juveniles, 19 males, 18 females). The faecal samples were stored in alcohol until analysis. In order to eliminate the effect of fragment size, each faecal sample was ground in a mortar and then passed through 0.8 and 0.4 mm sieves. The 0.4 and 0.8 fraction was dispersed in 50% glycerine solution. Three microscope slides were prepared for each sample and examined at $100\times$ and/or $200\times$ magnification under an optical microscope by two observers. Three horizontal traverses (2×60 mm) were scanned with 2 mm between traverses. The fragments in each traverse were identified to species level by comparing them to the reference collection, using epidermal characteristics (shape, size and arrangement of epidermal cells, stomata and hairs) and counted. The three slides results for each sample were pooled. The total fraction of unknown fragments remains low ($7\pm 3.5\%$).

2.4. Statistics

Due to small sample sizes and to the absence of normality in the distribution of the data, we used Kruskal Wallis tests to compare the numbers of plant species and the percentages of plant cover under and out of the jujube plant communities.

The data collected from faecal analysis were explored in two separate ways (1) the frequency of occurrence which was the proportion of faecal samples containing a given plant species and (2) the percent of each plant species in the diet. The proportion (p_i) of a given plant species 'i' in tortoise's faeces was calculated as the percentage of the number of epidermis fragment for each plant species relative to the total of the number of epidermis fragment we counted. Then, we calculated the mean p_i in tortoise's faeces for males, females and juveniles. Similarly, taxonomic availability in the study site (q_i) was calculated as the percentage of plant cover for each plant species relative to the total percentage of plant cover for each quadrat. Then, we calculated the mean q_i for the 43 quadrats we collected, and separately the mean q_i for quadrat under Jujube and quadrat out of Jujube. We used Hunter's index ($H_i = p_i/q_i$, Hunter, 1962) to distinguish preferred ($H_i > 1$) from avoided plant species ($H_i < 1$). Preference or avoidance were considered as statistically significant if the 95% confidence limits of mean p_i did not overlap the confidence limits of mean q_i . This statistical procedure was preferred to a simple contingency table with χ^2 tests because it takes into account the spatial heterogeneity of the vegetation and inter individual diet heterogeneity (Lagarde et al., 2003). The proportional similarity index ($PSI = 1 - 0.5 \sum_i |p_i - q_i|$) of Feisinger et al. (1981) was used to estimate to what extent *T. g. graeca* is a specialized or generalized diet. PSI can range from 1 when

animals consume any plant at their disposal to minimal ' q_i ' when they narrowly select their food (Feisinger et al., 1981).

Mean values were given with their standard deviation, and differences were considered statistically significant at $P < 0.05$. All statistical tests were performed using Statistica 5.1.

3. Results

3.1. Spatial variations in plant community

We identified 88 different plant species in our study area. The vegetation structure strongly diverged between quadrats performed under (UJ) and out of the Jujube (OJ). The mean plant species number and the total percent of recovery were significantly higher in UJ quadrats (mean = 90.2%) than in OJ quadrats (mean = 36.7%, Table 1). Species assemblage differed strongly between the both plant communities (Table 2).

3.2. Diet composition and food selection

We identified 34 different plant species in *T. graeca* diet. Because of the absence of significant differences between males, females and juveniles faecal samples, the results were pooled for subsequent analysis. Some invertebrate (insects and molluscs) fragments were found in 21% of faeces analysed.

Hunter's index calculation (Table 3) and the comparisons of the 95% confidence limits of mean p_i and of mean q_i (Table 2) suggest that eight plant species were significantly preferred and seven plant species significantly avoided (Table 2). The five main plant species recovered in faeces (*Leontodon saxatilis* (Asteraceae); *Malva parviflora* (Malvaceae); *Astragalus cruciatus*, *Medicago hispida* and *Lotus arenarius* (Fabaceae) represented about 70% of the identified material (Table 2). *L. saxatilis* seemed to be the most commonly consumed item (28%) followed by *M. parviflora* and *A. cruciatus* (15% each); these three species were regularly recorded (in more than 75% of the faeces analysed). *T. g. graeca* showed marked preferences for these five species, *Scorpiurus sulcata* was also a very common food item (Tables 2 and 3).

Table 1
Comparison between plant communities under and out of the jujube

	Under jujube	Out of jujube	K–W-test
Number of plant species	17.1 ± 7.3 (7–31)	10.9 ± 3.5 (4–19)	5.98; P = 0.01
Percentage of plant cover	90.2 ± 24.1 (48.8–112.0)	36.7 ± 17.1 (10.9–90.2)	16.98; P < 0.001

The mean number of plant species and the mean percentage of vegetation cover were given ± SD with min-max in parentheses. Kruskal–Wallis statistics were given in boldface when significant differences were detected between plant communities.

Table 2
Mean (\pm SD) plant availability (% of total vegetation cover) in the study site and percentage of species in the diet of *Testudo graeca graeca* during the spring 2002

Taxon	% of plant cover			Faecal samples		
	Total plant cover	Under jujube	Out Jujube	Mean number of fragments	Occurrence	% of occurrence
Aizoaceae						
<i>Aizoon hispanicum</i>	1.15 \pm 6.80(-0.89–3.18)	1.02 \pm 2.47(-0.06–2.63)	0.08 \pm 0.42(-0.06–0.23)	0.03 \pm 0.12(0.00–0.07)	3	6.8
<i>Aizoon canariense</i>	26.11 \pm 27.92(17.76–34.45)	2.53 \pm 4.05(-0.12–5.17)	32.32 \pm 28.78(22.65–42.00)	0	0	0
Alsinoideae						
<i>Spergularia flaccida</i>	3.53 \pm 7.06(1.42–5.64)	8.80 \pm 10.25(2.11–15.50)	1.75 \pm 3.00(0.74–2.76)	0.10 \pm 0.41(-0.02–0.22)	5	11.4
Caryophyllaceae						
<i>Fagopyra cretica</i>	1.06 \pm 4.00(-0.13–2.26)	5.25 \pm 9.34(-0.85–11.35)	0	0.03 \pm 0.20(-0.03–0.09)	1	2.3
<i>Spergularia marginata</i>	2.75 \pm 7.10(0.63–4.87)	2.13 \pm 5.12(-1.21–5.48)	3.07 \pm 8.05(0.36–5.77)	7.30 \pm 11.26(3.97–10.63)	36	81.8
Chenopodiaceae						
<i>Chenopodium murale</i>	0.27 \pm 0.76(0.05–0.50)	1.77 \pm 2.49(0.14–3.39)	0.02 \pm 0.12(-0.02–0.06)	0.02 \pm 0.11(-0.02–0.05)	1	2.3
Asteraceae						
<i>Anacyclus radiatus</i>	1.51 \pm 2.73(0.69–2.33)	8.05 \pm 11.76(0.37–15.73)	0.74 \pm 0.99(0.41–1.08)	0	0	0
<i>Calendula algeriensis</i>	0.38 \pm 1.21(0.02–0.75)	1.52 \pm 2.37(-0.02–3.07)	0.09 \pm 0.38(-0.03–0.22)	0.05 \pm 0.34(-0.05–0.15)	1	2.3
<i>Leontodon saxatilis</i>	0.99 \pm 2.56(0.23–1.76)	1.14 \pm 3.42(-1.09–3.37)	0.97 \pm 2.31(0.19–1.74)	27.91 \pm 18.37(22.48–33.33)	42	95.5
<i>Scolymus hispanicus</i>	0.65 \pm 1.91(0.08–1.22)	2.31 \pm 3.37(0.10–4.51)	0.23 \pm 1.13(-0.15–0.61)	0	0	0
<i>Asteraceae sp</i>	0.46 \pm 2.05(-0.15–1.07)	2.19 \pm 4.21(-0.56–4.94)	0	0	0	0
Convulvulaceae						
<i>Convolvulus althaeoides</i>	0.22 \pm 1.13(-0.12–0.56)	1.08 \pm 2.41(-0.49–2.66)	0	0	0	0
Brassicaceae						
<i>Biscutella didyma</i>	0.38 \pm 1.72(-0.13–0.89)	0.81 \pm 2.06(-0.54–2.15)	0.03 \pm 0.15(-0.02–0.08)	1.21 \pm 4.37(-0.08–2.50)	14	31.8
<i>Sinapsis arvensis</i>	0.37 \pm 1.57(-0.01–0.84)	2.66 \pm 4.83(-0.50–5.81)	0	0	0	0
<i>Notoceras bicorne</i>	1.14 \pm 1.56(0.67–1.60)	1.76 \pm 2.04(0.42–3.09)	1.12 \pm 1.50(0.62–1.63)	2.92 \pm 4.19(1.69–4.16)	32	72.7
Geraniaceae						
<i>Erodium guttatum</i>	0.27 \pm 1.16(-0.08–0.62)	1.25 \pm 2.49(-0.38–2.88)	0.02 \pm 0.13(-0.02–0.07)	0.01 \pm 0.06(-0.01–0.03)	1	2.3
Fabaceae						
<i>Medicago hispida</i>	0.05 \pm 0.20(-0.01–0.11)	0.11 \pm 0.25(-0.05–0.27)	0.04 \pm 0.22(-0.04–0.11)	5.75 \pm 9.85(2.84–8.66)	36	81.8
<i>Medicago laciniata</i>	12.52 \pm 14.59(8.16–16.88)	2.61 \pm 3.30(0.46–4.77)	14.29 \pm 16.70(8.68–19.90)	1.27 \pm 5.33(-0.30–2.85)	5	11.4

Table 2 (continued)

Taxon	% of plant cover		Out Jujube		Faecal samples		% of occurrence
	Total plant cover	Under jujube	Out Jujube	Mean number of fragments	Occurrence		
<i>Astragalus cruciatus</i>	0.02 ± 0.11 (-0.01-0.06)	0.08 ± 0.24 (-0.08-0.23)	0.01 ± 0.05 (-0.01-0.03)	14.37 ± 18.02 (9.04-19.70)	34	77.3	
<i>Lotus orenarius</i>	0.09 ± 0.36 (-0.02-0.20)	0.00	0.12 ± 0.42 (-0.02-0.27)	3.57 ± 5.96 (1.81-5.33)	25	56.8	
<i>Scorpiurus sulcata</i>	0.06 ± 0.29 (-0.03-0.14)	0.00	0.07 ± 0.32 (-0.04-0.18)	2.37 ± 5.58 (0.72-4.01)	18	40.9	
Malvaceae							
<i>Malva parviflora</i>	1.09 ± 4.62 (-0.29-2.47)	2.34 ± 2.74 (0.55-4.13)	0.96 ± 5.15 (-0.77-2.69)	15.22 ± 14.79 (10.85-19.58)	38	86.4	
Ombellifereae							
<i>Eryngium liticifolium</i>	20.21 ± 16.68 (15.23-25.20)	7.83 ± 6.68 (3.46-12.19)	23.33 ± 18.16 (17.22-29.43)	0	0	0	
<i>Anmi majus</i>	0.40 ± 1.61(-0.08-0.88)	1.92 ± 3.22(-0.18-4.02)	0	0.27 ± 0.64 (0.08-0.45)	12	27.3	
Plantaginaceae							
<i>Plantago amplexicaulis</i>	0.15 ± 0.64 (-0.04-0.34)	1.24 ± 2.62(-0.47-2.95)	0	2.82 ± 4.84 (1.39-4.25)	28	63.6	
Polygonaceae							
<i>Emex spinosus</i>	7.20 ± 10.30 (4.12-10.28)	2.92 ± 3.74(0.47-5.36)	8.93 ± 11.29 (5.13-12.73)	0.40 ± 1.20 (0.05-0.76)	11	25.0	
Scrophulariaceae							
<i>Linaria sagittata</i>	0.49 ± 3.24(-0.47-1.46)	2.36 ± 7.08(-2.26-6.98)	0	0	0	0	
<i>Linaria bipartita</i>	0.40 ± 0.91(0.13-0.67)	0	0.54 ± 1.08(0.18-0.91)	0.11 ± 0.76(-0.11-0.34)	1	2.3	
Solanaceae							
<i>Solanum nigrum</i>	0.58 ± 3.77(-0.55-1.70)	4.27 ± 12.82(-4.10-12.65)	0	0	0	0	
Poaceae							
<i>Festuca sp.</i>	3.19 ± 5.84(1.45-4.94)	0.06 ± 0.13 (-0.03-0.15)	3.75 ± 5.29 (1.98-5.53)	1.01 ± 2.34 (0.32-1.70)	12	27.3	
<i>Brachypodium distachyum</i>	2.36 ± 6.38(0.46-4.27)	10.99 ± 13.36(2.27-19.72)	0.14 ± 0.46 (-0.01-0.30)	3.52 ± 3.86 (2.38-4.66)	34	77.3	
<i>Cynodon dactylon</i>	1.11 ± 7.26(-1.06-3.28)	0	0.26 ± 1.52(-0.25-0.77)	0	0	0	
<i>Stipa retorta</i>	0.01 ± 6.62(-0.22-3.74)	7.05 ± 13.49(-1.76-15.87)	0.13 ± 0.38(0.00-0.26)	0	0	0	
Liliaceae							
<i>Asphodelus tenuifolius</i>	2.34 ± 5.12(0.81-3.87)	1.34 ± 1.57(0.31-2.37)	2.56 ± 5.56(0.69-4.43)	0.79 ± 1.84(0.25-1.33)	19	43.2	
Others	4.74 ± 5.99(2.95-6.53)	10.61 ± 9.57(4.36-16.86)	4.42 ± 4.21(3.00-5.83)	6.76 ± 3.50(5.72-7.79)	-	-	

Ninety-five per cent confidence intervals are indicated in brackets. Occurrence represents the number of faecal samples containing the item. The availability for each plant species in the study site was given in boldface when a significant difference was detected with the percentage of species in the diet. Data are based on 43 quadrats analyzed for plant availability (34 out of the Jujube and 9 under the Jujube) and 44 faecal samples of wild foraging tortoises.

Table 3

Hunter's index calculated for different plant species observed in the *Testudo g. graeca* diet in the Central Jbilets (Morocco)

Taxon	Hunter's index		
	Total plant cover	Under Jujube	Out Jujube
Aizoaceae			
<i>Aizoon hispanicum</i>	0.03	0.03	0.38
<i>Aizoon canariense</i>	0	0	0
Alsinoideae			
<i>Spergula flaccida</i>	0.03	0.01	0.06
Caryophyllaceae			
<i>Fagonia cretica</i>	0.03	0.01	—
<i>Spergularia marginata</i>	2.66	3.43	2.38
Chenopodiaceae			
<i>Chenopodium murale</i>	0.07	0.01	1.00
Asteraceae			
<i>Anacyclus radiatus</i>	0	0	0
<i>Calendula algeriensis</i>	0.13	0.03	0.56
<i>Leontodon saxatilis</i>	28.08	24.48	28.77
<i>Scolymus hispanicus</i>	0	0	0
<i>Asteraceaea sp</i>	0	0	—
Convolvulaceae			
<i>Convolvulus althaeoides</i>	0	0	—
Brassicaceae			
<i>Biscutella didyma</i>	3.19	1.49	40.33
<i>Sinapsis arvensis</i>	0	0	—
<i>Notoceras bicorne</i>	2.57	1.66	2.61
Geraniaceae			
<i>Erodium guttatum</i>	0.04	0.01	0.50
Fabaceae			
<i>Medicago hispida</i>	155.0	52.27	143.75
<i>Medicago laciniata</i>	0.10	0.49	0.09
<i>Astragalus cruciatus</i>	718.50	179.63	1437.0
<i>Lotus arenarius</i>	39.67	—	29.75
<i>Scorpiurus sulcata</i>	39.50	—	33.86
Malvaceae			
<i>Malva parviflora</i>	13.96	6.50	15.85
Ombellifereae			
<i>Eryngium ilicifolium</i>	0	0	0
<i>Ammi majus</i>	0.68	0.14	—
Plantaginaceae			
<i>Plantago amplexicaulis</i>	18.80	2.27	—
Polygonaceae			
<i>Emex spinosus</i>	0.06	0.14	0.04
Scrophulariaceae			
<i>Linaria sagittata</i>	0	0	—
<i>Linaria bipartita</i>	0.14	0.03	0.61
Solanaceae			
<i>Solanum nigrum</i>	0	0	—

Table 3 (continued)

Taxon	Hunter's index		
	Total plant cover	Under Jujube	Out Jujube
Poaceae			
<i>Festuca sp.</i>	0.32	16.83	0.27
<i>Brachypodium distachyum</i>	1.49	0.32	25.14
<i>Cynodon dactylon</i>	0	—	0
<i>Stipa retorta</i>	0	0	0
Liliaceae			
<i>Asphodelus tenuifolius</i>	0.34	0.59	0.31

However, the tortoises actively avoided *Spergula flaccida*, *Eryngium ilicifolium*, and *Anacyclus radiatus*. For nine plant species (*Aizoon canariense*, *Ammi majus*, *Brachypodium distachyum*, *Chenopodium murale*, *Emex spinosus*, *M. laciniata*, *Notoceras bicorne*, *Plantago amplexicaulis*, *Scolymus hispanicus*) conventional significance for selectivity was reached in one plant community solely (Table 2). Finally, a more puzzling result (inconsistent trend under and out of the Jujube) was obtained for a single species (*Festuca sp.*) (Table 2). It should be noted however that these eight plant species was only rarely found as a food item (0–3.5% of identified fragments).

The PSI was always less than 0.25 (0.20 when considering the overall plant community, and 0.24 and 0.17 when focussed on under the Jujube and out of the Jujube plant communities, respectively), indicating that *T. g. graeca* is a rather selective herbivore.

4. Discussion

4.1. Dietary

Our study provides the first quantitative data on the feeding ecology of *T. g. graeca* using faecal analysis method. The obtained data indicate clearly that *T. g. graeca* is essentially herbivorous, with 34 different species identified in the faeces. This result agree with the general tendency founded in most Testudinidae species, though some occasionally take other material, including fungi, invertebrates, carrion, bones and soil (Pritchard, 1979; Moskovits and Bjorndal, 1990; Milton, 1992; Rall and Fairall, 1993; Bayley and Highfield, 1996). Such diversity may allow the Spur-thighed tortoise to reach an adequate balance of nutrients by ingesting a wide variety of plants (until 18 plant species in single faeces sample). Therefore, diet mixing in herbivores can greatly increase digestibility of dry matter, organic matter and energy sources (Bjorndal, 1991). In addition to these wild plant species, the diet of *T. g. graeca* also include a wide variety of cultivated plants (lucerne, potato leaves, tomatoes, lettuce, etc.) in the Agadir region (Bayley and Highfield, 1996). According

to these authors, the loss of vegetal cover and drought contributed to increase tortoise population densities around the intensively farmed areas and then changed their feeding habits by consuming the cultivated species. In Spain, [Andreu \(1987\)](#) recorded 88 plant species in the diet of Spur-thighed inhabiting the Doñana Park. This number of plant is high compared to the 34 plant species found in our study. This could be due to local bio-climatic conditions. Nevertheless, the major constituents of tortoise diets in the two area (Jbilet and Doñana) were essentially composed on Asteraceae, Fabaceae and Poaceae.

4.2. Is *T. g. graeca* a generalist or a specialist feeder?

Our results suggested that five plant species made up 70% of the diet (*L. saxatilis*, *A. cruciatus*, *M. parviflora*, *M. hispida*, *L. arenarius*). Surprisingly, these main plant items consumed correspond to relatively under represented species in plant community and other ones like *A. canariense* and *E. ilicifolium* are very abundant in the environment during the study period (represent 46% of the total vegetal cover) and highly avoided (0% in the diet). This suggest that Spur-thighed tortoise strongly select its food, and the PSI ([Feisinger et al., 1981](#)) confirmed this tendency (PSI < 0,25 in all the cases): *T. g. graeca* in the Jbilet during spring is rather a specialist herbivorous species. This strongly contrasts with other studies but one (*Gopherus agassizii*: [Jennings, 1993](#)), where Chelonian species are generally at midway between a specialist and a generalist herbivore (*G. polyphemus*: [MacDonald and Mushinsky, 1988](#), *Geochelone pardalis*: [Mason et al., 1999](#), *G. denticulata* and *G. carbonaria*: [Bjorndal, 1989](#); [Moskovits and Bjorndal, 1990](#), *T. horsfieldi*: [Lagarde et al., 2003](#)).

4.3. Why *T. g. graeca* select its food?

The major factors influencing feeding selectivity in herbivorous species seems to be principally the energetic, hydric and toxic contents of plants consumed (see [Pyke, 1984](#) for a review and [Moskovits and Bjorndal, 1990](#); [Mason et al., 1999](#); [Lagarde et al., 2003](#)). [Belovsky \(1981\)](#) and [Owen-Smith and Novellie \(1982\)](#) recognized that concentrations of specific nutrients in potential foods may play a significant role in determining diet choice. [Bjorndal \(1980\)](#) found that the herbivore sea turtles, *Chelonia mydas*, select a food with both high protein and low lignin content to increase digestibility. [Garner and Landers \(1981\)](#) predicted *G. polyphemus* diet selection according to nutritional compound. They considered the Fabaceae as the most important forbs in the diet both because of a high nutritional value. Another study on *G. agassizii* ([Jennings 1993](#)) showed similar results: four species of Fabaceae (*Lotus* and *Astragalus* species) were among the most highly preferred plant species and represented about half of the seasonal diet of the tortoises. In the present study too, within eight plant species significantly preferred by *T. g. graeca*, four are Fabaceae representing 27% of total diet. *E. ilicifolium* is probably unpalatable and hard to digest by herbivores ([Floret, 1988](#)) because of its thorny leaves (external defence mechanisms) and then highly avoided by tortoise. The present study showed that *T. g. graeca* consumes some plants (e.g. *M. parviflora*, *Solanum nigrum*,

A. majus, *Anagalis arvensis* and *C. murale*) which could be toxic for mammalian herbivores because of toxic contents (Halsey, 1998; Forshaw, 2000; Knight and Walter, 2002). The occurrence of potentially toxic items in the diet has been documented in other Chelonian species (*T. hermanni*: Longepierre and Grenot, 1999; *Eretmochelys imbricata*: Meylan, 1988; *T. graeca*: Andreu et al., 2000; *T. horsfieldi*: Lagarde et al., 2003) and has been interpreted as anti-helminthic strategy (Satorhelyi and Sreter, 1993; Longepierre and Grenot, 1999, Lagarde et al., 2003).

Despite the consumption of some plants species that could be toxic for mammalian herbivores, the present study indicated that in our study area, *T. g. graeca* feed mostly on plant species that are also appreciated by the domestic mammals (Floret, 1988). Then, competition for food between tortoises and domestic mammals could occur in overgrazing areas and in such a context, spiny shrubs may represent a crucial protection for tortoises feeding resources.

4.4. Where *T. g. graeca* forage?

For several (10 out 34) plant species, the significance of plant selection and/or the Hunter selectivity indexes differs when comparing the diet of the tortoises under and out of the Jujube shrubs. However, these problematic plant species represent only a marginal part of the diet of the tortoises (0–3.5% of the plant fragments identified). It is therefore difficult to consider that the inconsistency in some of our results reflects a real change in the feeding selectivity of the tortoises. More likely, the abundance of such plants strongly differs between the two plant communities and/or, *T. g. graeca* prefers to forage under Jujube, where the global feeding resources are more abundant and diversified.

4.5. Methodological bias and perspectives

We focussed our study on the spring period. It could be very useful to study the feeding preferences seasonal variations. But nevertheless, spring is the period of maximum activity level and then the period of maximum food intake.

Faecal analysis was classically considered as a limited method in describing diet composition quantitatively (Rice, 1970; Slater and Jones, 1971; McInnis et al., 1983; Gordon, 1995), due to differential digestion rates of plant species and the typically high proportion of unidentified cuticle remains occurring in the faeces. Nevertheless, in our results, the percentage of unidentified remains (6.6%) and the percentage of wrong recognition ($7 \pm 3.5\%$) were low. Of course, we cannot exclude problems of differential digestibility of plant items and we need additional analysis to confirm the major pattern we found: *T. g. graeca* seems a rather specialist herbivorous species.

Acknowledgements

Substantial financial support was provided by the Integrated Action number MA/02/54 (International Program for Scientific Collaboration between Morocco

and France) granted by the “Comité Mixte Inter Universitaire Franco-Marocain”. The “Conseil Général des Deux Sèvres” improved logistical support. We warmly thank Hussein Boujoudi and his family for his help in the field and Nathalie Bourgougnon for references on plant toxicity. Rex Cambag kindly lent its sophisticated field material.

References

- Andreu, A. C., 1987. Ecologia dinamica poblacional de la tortuga mora, *Testudo graeca*, en Doñana. Thesis doctoral, University of Seville.
- Andreu, A.C., Diaz-Paniagua, C., Keller, C., 2000. La tortuga mora (*Testudo graeca* L.) en Doñana. Asociaciou Herpetologica Espanola, Monografias de Herpetologia, vol. 5.
- Baumgartner, L.L., Martin, A.C., 1939. Plant histology as an aid in squirrel food-habit studies. *Journal of Wildlife Management* 3, 266–268.
- Bayley, J.R., Highfield, A.C., 1996. Observations on ecological changes threatening a population of *Testudo graeca graeca* in the Souss Valley, Southern Morocco. *Chelonian Conservation and Biology* 2, 36–42.
- Belovsky, G.E., 1981. Food plant selection by a generalist herbivore : the moose. *Ecology* 62, 1020–1030.
- Bjorndal, K.A., 1980. Nutrition and grazing behavior of the Green Turtle *Chelonia mydas*. *Marine Biology* 56, 147–154.
- Bjorndal, K.A., 1989. Flexibility of digestive responses in two generalist herbivores, the tortoises *Geochelone carbonaria* and *Geochelone denticulata*. *Oecologia* 78, 317–321.
- Bjorndal, K.A., 1991. Diet mixing: non-additive interactions of diet items in an omnivorous freshwater turtle. *Ecology* 72, 1234–1241.
- Bonnet, X., Lagarde, F., Henen, B.T., Corbin, J., Nagy, K.A., Naulleau, G., Balhoul, K., Chastel, O., Legrand, A., Cambag, R., 2001. Sexual dimorphism in steppe tortoises (*Testudo horsfieldi*): influence of the environment and sexual selection on body shape and mobility. *Biological Journal of the Linnean Society* 72, 357–372.
- Chapuis, J.L., 1980. Méthodes d'étude du régime alimentaire de Lapin de garenne *Orytolagus cuniculus* (L.) par l'analyse microscopique des fèces. *Revue d'Ecologie-La Terre et la Vie* 34, 159–198.
- Chapuis, J.L., Didillon, M.C., 1987. Méthodes d'étude du régime alimentaire des Galliformes. Issue Series Title: Gibier Faune Sauvage 4, 295–320.
- Cobo, M., Andreu, A.C., 1988. Seed consumption and dispersal by the spur-thighed tortoise, *Testudo graeca graeca*. *Oikos* 51, 267–273.
- El Mouden, E.H., Znari, M., Brown, R., 1999. Skeletochronology and mark-recapture assessments of growth in the North African agamid lizard (*Agama impalearis*). *Journal of Zoology of London* 249, 455–461.
- El Mouden, E. H., Slimani, T., Ben Kaddour, K., 2001. Croissance et dimorphisme sexuel chez la Tortue Mauresque (*Testudo graeca graeca* L. 1758). *Chelonii* 3, Proceeding International Congress on Testudo Genus, March 7–10.
- Emberger, L., 1933. Nouvelle contribution à l'étude de la classification des groupements végétaux. *Revue Générale de Botanique* 45, 473–486.
- Feisinger, P.E., Spears, E., Poole, R.W., 1981. A simple measure of niche breadth. *Ecology* 62, 27–32.
- Floret, C., 1988. Méthodes de mesure de la végétation pastorale. *Pastoralisme et développement*, 24 mai-9 juillet, IAV. Hassan II, Rabat, Maroc, IAM, Montpellier, France.
- Forshaw, D., 2000. “Redgut” a cause of deaths in sheep grazing lucerne and other legumes. *Animal Health, Animal Disease Surveillance Newsletter, The Agriculture Protection Program*, December 2.
- Garcia-Gonzalez, R., 1984. L'emploi des épidermes végétaux dans la détermination du régime alimentaire de l'Isard dans les Pyrénées occidentales. *Ecologie des milieux montagnards et de haute altitude. Documents d'Ecologie Pyrénéenne III-IV*, 307–313.

- Garner, J., Landers, J., 1981. Foods and habitat of the gopher tortoise in southwestern Georgia. *Proceeding of the Annual Conference on Southeast Association of Fisheries and Wildlife Agencies*, vol. 35, pp. 120–134.
- Gordon, I.J., 1995. Animal-based techniques for grazing ecology research. *Small Ruminant Research* 16, 203–214.
- Halsey, L.A., 1998. Nitrate in forage cause cattle deaths: a common weed and uncommon circumstances. *Managing nutrition and forages to improve productivity and profitability. Florida Beef Cattle Short Course*, pp. 99–108.
- Hercus, B.H., 1960. Plant cuticle as an aid to determining the diet of grazing animals. *Proceedings of the Eighth International Grassland Congress*, pp. 443–447.
- Highfield, A.C., 1989. Notes on dietary constituents for herbivorous terrestrial chelonia and their effects on growth and development. *ASRA (UK) Journal* 3, 7–20.
- Highfield, A.C., 1990a. Preliminary Report on the Taxonomic, Biotypic and Conservation Status of the Land Tortoises of Tunisia. *Tortoise Trust, London* 14pp.
- Highfield, A.C., 1990b. Tortoise of North Africa: taxonomy nomenclature, phylogeny and evolution with notes on field studies in Tunisia. *Journal of Chelonian Herpetology* 1, 1–56.
- Highfield, A.C., 1994. The Illegal Trade to Tourist in Morocco of Souvenir Products made from Endangered Tortoises. *Tortoise Trust, London*.
- Hunter, R.F., 1962. Hill sheep and their pasture : a study of sheep grazing in south East Scotland. *Journal of Ecology* 50, 551.
- Jennings, W. B., 1993. Foraging ecology of the desert tortoise (*Gopherus agassizii*) in the western Mojave desert. M.S. Thesis, University of Texas at Arlington.
- Knight, A.P., Walter, R.G., 2002. Plant causing sudden death. In: Knight, A.P., Walter, R.G. (Eds.), *Plant Poisoning of Animals in North America*. Teton Newmedia, Jackson WY, p. 41.
- Lagarde, F., Bonnet, X., Nagy, K., Henen, B., Corbin, J., Naulleau, G., 2002. A short spring before a long jump: the ecological challenge to the steppe tortoise (*Testudo horsfieldi*). *Canadian Journal of Zoology* 80, 493–502.
- Lagarde, F., Bonnet, X., Corbin, J., Henen, B., Nagy, K., Mardonov, B., Naulleau, G., 2003. Foraging behaviour and diet of an ectothermic herbivore: *Testudo horsfieldi*. *Ecography* 26, 236–242.
- Lambert, M.R.K., 1969. Tortoise drain in Morocco. *Oryx* 10, 161–166.
- Lambert, M.R.K., 1981. Temperature, activity and field sighting in the Mediterranean spur-thighed or common garden tortoise *Testudo graeca* L. *Biological Conservation* 21, 39–54.
- Lambert, M.R.K., 1983. Some factors influencing the Moroccan distribution of the western Mediterranean spur-thighed tortoise. *Testudo graeca graeca* L. and those predclufing its survival in NW Europe. *Zoological Journal of the Linnean Society* 79, 149–179.
- Le Hou erou, H., 1989. Classification  eoclimatique des zones arides (s.l.) de l’Afrique du Nord. *Ecologia Mediterranea* XV, 95–144.
- Loehr, V.J.T., 2002. Diet of the Namaqualand speckled padloper, *Homopus signatus signatus*, in early spring. *African Journal of Herpetology* 51, 47–55.
- Longepierre, S., Grenot, C., 1999. Some effects of intestinal nematodes on the plant foraging behaviour of *Testudo hermani hermani* in the south of France. In: Miaud, C., Guyetant, G. (Eds.), *Current Studies in Herpetology. Le Bourget du Lac (SEH)*, pp. 277–284.
- Macdonald, L.A., Mushinsky, H.R., 1988. Foraging ecology of the Gopher tortoise, *Gopherus polyphemus*, in a Sandhill Habitat. *Herpetologica* 44, 345–353.
- Mason, M.C., Kerley, G.I.H., Weatherby, C.A., Branch, W.R., 1999. Leopard Tortoises (*Geochelone pardalis*) in Valley Bushveld, Eastern Cape, South Africa: specialist or generalist herbivores? *Chelonian Conservation and Biology* 3, 435–440.
- McInnis, M.L., Varva, M., Krueger, W.C., 1983. A comparison of four methods used to determine the diets of large herbivores. *Journal of Range Management* 36, 730–732.
- Meylan, A.B., 1988. Spongivory in hawksbill turtle: a diet of glass. *Science* 239, 393.
- Milton, S.J., 1992. Plant eaten and dispersed by adult leopard tortoises *Geochelone pardalis* (Reptilia: Chelonii) in the southern Karoo. *South African Journal of Zoology* 27, 45–49.

- Moskovits, D.K., Bjorndal, K.A., 1990. Diet and food preferences of the tortoises *Geochelone carbonaria* and *G. denticilata* in northwestern Brazil. *Herpetologica* 46, 207–218.
- Nègre, R., 1961. Petite flore des régions arides du Maroc occidental. Centre National de la Recherche Scientifique, Tome I, pp. 413.
- Nègre, R., 1962. Petite flore des régions arides du Maroc occidental. Centre National de la Recherche Scientifique, Tome II, pp. 566.
- Owen-Smith, N., Novellie, P., 1982. What should a clever ungulate eat? *American Naturalist* 119, 151–178.
- Pritchard, P.C.H., 1979. *Encyclopedia of Turtles*. TFH Publ., Neptune, NJ 895 pp.
- Pyke, G.H., 1984. Optimal foraging theory : a critical review. *Annual Review of Ecology and Systematics* 15, 523–575.
- Quezel, P., Santa, S., 1962. Nouvelle flore de l'Algérie et des régions désertiques méridionales. Centre National de la Recherche Scientifique, Tome I, Paris, pp.1–565.
- Quezel, P., Santa, S., 1963. Nouvelle flore de l'Algérie et des régions désertiques méridionales. Centre National de la Recherche Scientifique, Tome II, Paris, pp. 571–1170.
- Rall, M., Fairall, N., 1993. Diets and food preferences of two South African tortoises *Geochelone pardalis* and *Psammodromus ocellifer*. *South African Journal of Wildlife Research* 23, 63–70.
- Rice, R. W., 1970. Stomach content analysis: a comparison of the rumen vs. oesophageal techniques. In: *Range and Wildlife Habitat Evaluation—A Research Symposium*. United States Department of Agriculture and Forestry Services. Miscellaneous Publications 1147, 127–132.
- Satorhelyi, T., Sreter, T., 1993. Studies on internal parasites of tortoises. Issue Series Title: *Parasitologia Hungarica* 26, 51–55.
- Slater, J., Jones, R.J., 1971. Estimation of diets selected by grazing animals from microscopic analysis of the faeces—a warning. *Journal of the Australian Institute of Agricultural Science* 37, 238–240.
- Slimani, T., El Mouden, E. H., Ben Kaddour, K., 2001. Structure et dynamique d'une population de *Testudo graeca graeca* L. 1758 dans les Jbilet Centrales, Maroc, *Chelonii* 3, Proceeding International Congress on Testudo Genus, March 7–10, Gonfaron, France.
- Sparks, D.R., Malechek, J.C., 1968. Estimating percentage dry weight in diets using a microscopic technique. *Journal of Range Management* 32, 275–279.
- Stewart, D.R.M., 1967. Analysis of plant epidermis in faeces: a technique for studying the food preferences of grazing herbivores. *Journal of Applied Ecology* 4, 83–111.
- Stewart, D.R.M., 1970. Survival during digestion of epidermis from plants eaten by ungulates. *Revue de Zoologie et de Botanique Africaines* 83, 343–348.
- Varva, M.V., Holecek, J.L., 1980. Factors influencing microhistological analysis of herbivore diets. *Journal of Range Management* 33, 371–374.
- Varva, M., Rice, R.W., Hansen, R.M., 1978. A comparison of esophageal fistula and fecal materials to determine deer diets. *Journal of Range Management* 31, 11–13.
- Znari, M., El Mouden, E., Francillon-Vieillot, H., 2002. Long-term variation in reproductive traits of Bibron's Agama, *Agama impalearis*, in Western Morocco. *African Journal of Herpetology* 51, 57–68.