

# Reproductive performance and diving behaviour share a common sea-ice concentration optimum in Adélie penguins (*Pygoscelis adeliae*)

Camille Le Guen<sup>1</sup>  | Akiko Kato<sup>1</sup> | Ben Raymond<sup>2</sup> | Christophe Barbraud<sup>1</sup> |  
 Michaël Beaulieu<sup>3,4</sup> | Charles-André Bost<sup>1</sup> | Karine Delord<sup>1</sup> | Andrew J. J. MacIntosh<sup>5</sup> |  
 Xavier Meyer<sup>6</sup> | Thierry Raclot<sup>6</sup> | Michael Sumner<sup>2</sup> | Akinori Takahashi<sup>7,8</sup> |  
 Jean-Baptiste Thiebot<sup>7</sup> | Yan Ropert-Coudert<sup>1</sup>

<sup>1</sup>Centre d'Etudes Biologiques de Chizé, UMR7372 CNRS-Université La Rochelle, Villiers en Bois, France

<sup>2</sup>Australian Antarctic Division, Department of the Environment, Australian Government, Kingston, Tasmania, Australia

<sup>3</sup>Zoological Institute & Museum, University of Greifswald, Greifswald, Germany

<sup>4</sup>German Oceanographic Museum, Stralsund, Germany

<sup>5</sup>Kyoto University Primate Research Institute, Inuyama, Japan

<sup>6</sup>CNRS, Institut Pluridisciplinaire Hubert Curien UMR7178, Université de Strasbourg, Strasbourg, France

<sup>7</sup>National Institute of Polar Research, Tachikawa, Tokyo, Japan

<sup>8</sup>Department of Polar Science, SOKENDAI (The Graduate University for Advanced Studies), Tokyo, Japan

## Correspondence

Camille Le Guen, Pelagic Ecology Research Group - School of Biology, University of St. Andrews, St-Andrews, Fife, UK.  
 Email: cam.leguen@gmail.com

## Funding information

The Japanese Government Mombukagakusho; The Japanese Society for the Promotion of Science; Institut Polaire Français Paul Emile Victor; World Wildlife Fund; Zone Atelier Antarctique et Subantarctique; SENSEI project from the BNP Paribas Climate Foundation

## Abstract

The Southern Ocean is currently experiencing major environmental changes, including in sea-ice cover. Such changes strongly influence ecosystem structure and functioning and affect the survival and reproduction of predators such as seabirds. These effects are likely mediated by reduced availability of food resources. As such, seabirds are reliable eco-indicators of environmental conditions in the Antarctic region. Here, based on 9 years of sea-ice data, we found that the breeding success of Adélie penguins (*Pygoscelis adeliae*) reaches a peak at intermediate sea-ice cover (ca. 20%). We further examined the effects of sea-ice conditions on the foraging activity of penguins, measured at multiple scales from individual dives to foraging trips. Analysis of temporal organisation of dives, including fractal and bout analyses, revealed an increasingly consistent behaviour during years with extensive sea-ice cover. The relationship between several dive parameters and sea-ice cover in the foraging area appears to be quadratic. In years of low and high sea-ice cover, individuals adjusted their diving effort by generally diving deeper, more frequently and by resting at the surface between dives for shorter periods of time than in years with intermediate sea-ice cover. Our study therefore suggests that sea-ice cover is likely to affect the reproductive performance of Adélie penguins through its effects on foraging behaviour, as breeding success and most diving parameters share a common optimum. Some years, however, deviated from this general trend, suggesting that other factors (e.g. precipitation during the breeding season) might sometimes become preponderant over the sea-ice effects on breeding and foraging performance. Our study highlights the value of monitoring fitness parameters and individual behaviour concomitantly over the long-term to better characterize optimal environmental conditions and potential resilience of wildlife. Such an approach is crucial if we want to anticipate the effects of environmental change on Antarctic penguin populations.

## KEYWORDS

Antarctic, breeding success, diving activity, foraging strategies, *Pygoscelis adeliae*, sea ice

## 1 | INTRODUCTION

Marine ecosystems are experiencing various disturbances (Richardson & Polocsanska, 2008), such as climate change (Doney et al., 2012; IPCC, 2007), overfishing (Jennings & Kaiser, 1998) and invasion of exotic species (Elton, 1958; Katsanevakis et al., 2014). Determining how and to what extent organisms are able to cope with environmental changes is fundamental to understand and protect marine ecosystems. This is especially needed in the Antarctic, parts of which have shown rapid change in recent decades (Clarke & Harris, 2003). In these marine ecosystems, the seasonal and interannual sea-ice dynamics are the main drivers of biogeochemical cycles (Sedwick & DiTullio, 1997; Wang, Bailey, Lindsay, Moore, & Holland, 2014). Seasonal changes in sea-ice cover alter nutrient cycling and induce seasonal variation in nutrient availability (Wang et al., 2014). These seasonal changes also control phytoplankton blooms, which occur after sea ice locally breaks up (Smith & Nelson, 1985; Wang et al., 2014). Importantly, sea ice provides a favourable habitat for microalgae and bacteria (Thomas & Dieckmann, 2010) that are able to cope with variable salinity conditions and to photosynthesize even under low light conditions (Knox, 2006). In turn, some krill species (e.g. the Antarctic krill, *Euphausia superba*) feed on these microalgae (Brierley, 2002), and predators can take advantage of this association to feed on concentrated prey (Knox, 2006). Sea ice therefore shapes the structure and functioning of the Antarctic ecosystem, from grazers to the highest trophic levels, and changes in its dynamics are likely to disrupt this interplay.

At a time when climatic conditions are changing rapidly, the Antarctic ecosystem requires close monitoring. However, monitoring an entire ecosystem is logistically challenging. To circumvent this difficulty, meso- and top-predators like seabirds and marine mammals can be used as eco-indicators of changes in their respective ecosystems (Boyd & Murray, 2001; Frederiksen, Mavor, & Wanless, 2007; Furness & Camphuysen, 1997), as they integrate and amplify the perturbations occurring at lower trophic levels (Hindell, Bradshaw, Harcourt, & Guinet, 2003). Seabirds are widely used as eco-indicators because they are sensitive to environmental variability in marine ecosystems (Bost et al., 2009; Croxall, Trathan, & Murphy, 2002; Smith et al., 1999). This logic formed the basis of the Conservation of Antarctic Marine Living Resources (CCAMLR) Ecosystem Monitoring Program (CEMP; CCAMLR, 2014). In addition, seabirds are central-place foragers (Orians & Pearson, 1979), meaning that individuals return regularly to their nesting site during the breeding season, which makes them easily accessible to researchers (Piatt, Sydeman, & Wiese, 2007). In Antarctica, changes in sea-ice cover are major determinants of seabird survival and reproduction (Barbraud & Weimerskirch, 2003; Croxall et al., 2002; Jenouvrier et al., 2014). Sea ice can be a primary factor affecting the foraging patterns of breeding adults, the provisioning of chicks, the breeding success, and ultimately the population dynamics (Fraser, Trivelpiece, Ainley, & Trivelpiece, 1992; Kato, Ropert-Coudert, & Naito, 2002). This highlights the need to investigate the relationship between sea-ice conditions and seabird foraging behaviour if we are to use such species as

indicators of changes in this polar ecosystem. Developments in bio-logging technology have greatly facilitated advances in our state of knowledge about seabird foraging behaviour and indicators of prey availability (Bost et al., 2009). Indeed, seabirds can now be tracked at increasingly finer scale and higher resolution, enhancing their value as eco-indicating species (Ropert-Coudert, Kato, Gremillet, & Grenner, 2012; Ropert-Coudert & Wilson, 2005).

Among seabirds, penguins represent up to 90% of the total avian biomass in the Southern Ocean (Knox, 2006). Adélie penguins (*Pygoscelis adeliae*) are predators that need to dive to forage, as they mostly feed on Antarctic krill (*E. superba*), ice krill (*E. crystallophias*), Antarctic silverfish (*Pleuragramma antarcticum*; Cherel, 2008; Croxall & Lishman, 1987; Libertelli, Coria, & Marateo, 2003; Volkman, Presler, & Trivelpiece, 1980) and to a lower extent on jellyfish (Thiebot et al., 2016). With its typically ice-associated prey and its circumpolar distribution, the Adélie penguin is a relevant candidate as an indicator of the state of sea-ice ecosystem (Ainley, 2002; CCAMLR, 2014; Woehler & Johnstone, 1991). Few studies have previously investigated the relationship between Adélie penguin diving behaviour and sea-ice dynamics. Among them, Emmerson and Southwell (2008) investigated the breeding and foraging performance of Adélie penguins in relation to annual sea-ice conditions. However, detailed information on their foraging behaviour, especially regarding the diving activity, which is required to genuinely link such responses to sea-ice conditions, remains rare, as does any understanding of the associated mechanisms. Other studies have typically compared only two or three penguin breeding sites for a given year (Watanuki, Kato, Naito, Robertson, & Robinson, 1997; Watanuki et al., 2002) and/or seasons with contrasting sea-ice conditions at a given site (Ainley et al., 1998; Beaulieu et al., 2010; Kato, Watanuki, & Naito, 2003), thereby limiting the generality of their results. Some studies have also investigated the relationship between sea-ice concentration and foraging efficiency (Lescroël, Ballard, Gremillet, Authier, & Ainley, 2014) or breeding success and survival (Dugger, Ballard, Ainley, Lyver, & Schine, 2014; Lescroël, Dugger, Ballard, & Ainley, 2009) over the long-term, but none have included analyses of fine-scale diving activity. However, all these studies converged towards the conclusion that sea ice plays a fundamental role in determining foraging strategies, and it was suggested that there may be an optimal range of sea-ice cover during the breeding season that leads to maximal foraging and breeding success (Ainley, 2002; Ballard, Dugger, Nur, & Ainley, 2010; Barbraud, Delord, & Weimerskirch, 2015; Smith et al., 1999). In this context, long-term monitoring of the foraging activity and breeding performance of Adélie penguins covering a wide array of sea-ice conditions is required to better understand and predict how these predators will respond to the forthcoming environmental changes in the Southern Ocean, especially given that the total extent of Antarctic sea-ice showed a counterintuitive, small increase of 1.5% per decade over the past 34 years, with strong regional differences around the Antarctic (Vaughan et al., 2013).

In this study, we investigate how changes in sea-ice characteristics at a given location affect the breeding success and diving activity of Adélie penguins. We aimed to test the robustness of the

conceptual model of Smith et al. (1999) and determine whether an optimal sea-ice concentration is indeed observable for this species not only in relation to breeding success but also to fine-scale metrics characterizing foraging activity, namely diving activity. Towards this end, we monitored the breeding success and diving behaviour, at multiple measurement scales, of chick-rearing Adélie penguins from a single colony in Adélie Land, East Antarctica, over nine austral summers with variable sea-ice conditions. The last aim of our study was to assess the relevance of Adélie penguins as eco-indicators of the Southern Ocean ecosystem, a key step towards addressing question 65 of the 1st Antarctic and Southern Ocean Science Horizon Scan of SCAR (Scientific Committee on Antarctic Research), i.e. “What will key marine species tell us about trophic interactions and their oceanographic drivers such as future shifts in frontal dynamics and stratification?” (CCAMLR, 2014; Kennicutt et al., 2014).

## 2 | MATERIALS AND METHODS

An Adélie penguin breeding colony located near the Dumont d'Urville station (66°40'S, 140°01'E), Adélie Land, Antarctica (Figure 1) was studied over nine nonconsecutive austral summers (October to March) between 1995 and 2014. To reduce variability due to differences in foraging strategies across the breeding cycle (Widmann et al., 2015), we focused on the guard stage (end of December - beginning of January), during which one parent guards the chicks on the nest while the other is foraging at sea to bring food back to its offspring. A total of 121 birds (with one foraging trip for each individual) were monitored (Supporting Information Table S1 for details).

### 2.1 | Sea-ice characteristics

We used satellite-derived passive microwave measurements of daily sea-ice concentration (SIC) (Cavalieri, Parkinson, Gloersen, & Zwally, 1996) to characterize the sea-ice conditions encountered by the studied individuals in every considered year. Other studies have shown that foraging trips of Adélie penguins at the Dumont d'Urville colony extend from 63.7°S to 66.6°S and from 134.7°E to 142.3°E across the breeding season, corresponding to an area of 119,389 km<sup>2</sup> or a foraging distance (maximum distance from the colony) of 392 km (Cottin et al., 2012; Widmann et al., 2015). Hence, we considered a slightly larger area from which we extracted sea-ice data to ensure complete cover of their foraging domain (62°–67°S and 134°–144°E; Figure 1). Sea-ice data were processed using R (R version 3.2.3, R Development Core Team, 2015) and the package “raster” (Hijmans et al., 2016) with a resolution of 25 km, which was the smallest common resolution for all years. Daily maps were created with a single value of SIC in each pixel of the raster (62°–67°S, 134°–144°E; Figure 1). SIC describes what percentage of a 25 × 25 km box is covered with ice, 0% being open water and 100% being full ice cover. For each day, the mean value of SIC was calculated over all pixels. The overall SIC value was calculated as the mean of the daily SIC values extracted within the area between the

1st of November (beginning of the season) and the 15th of January (end of diving data).

### 2.2 | Adélie penguin breeding success


Breeding success of Adélie penguins in Dumont d'Urville was monitored annually from 1995 to 2014 (see Jenouvrier, Barbraud, & Weimerskirch, 2006 and Barbraud et al., 2015 for further details). Breeding success is defined here as the ratio of the number of chicks counted in the area in early February (just before fledging) to the number of incubating pairs in late November (just after laying). A generalized additive model (GAM), which is a nonparametric smoothing regression technique, specified with a Gaussian error distribution, was fitted on the data available (20 years from 1995 to 2014) to investigate the relationship between breeding success and the overall SIC. The quadratic aspect of the relationship between these two variables was tested comparing a GAM with and without including a quadratic term (models compared using the Akaike Information Criterion, AIC). We used the “mgcv” package in R (version 3.2.3) to fit the GAM to our data (Wood, 2006), using a cubic regression spline. We used three knots as there were less than 30 observations, following the optimal amount of smoothing recommended by Zuur, Ieno, Walker, Saveliev, and Smith (2009).

### 2.3 | Adélie penguin foraging activity

Diving behaviour was recorded by miniature data loggers attached to the birds. Birds were captured while, or just before, leaving their nest for a foraging trip and equipped with data-loggers. These devices recorded time series of depth readings taken regularly at predetermined intervals (1 or 5 s; Supporting Information Table S1). Loggers were attached to the lower back of penguins using waterproof tape (Wilson & Wilson, 1989), except the LUL loggers used in 2014 which, given their small size, were attached to a leg band (Ratcliffe et al., 2014). After one or several foraging trips, birds were recaptured upon their return to the colony and the loggers were retrieved. Upon recovery, depth data were downloaded onto a computer and analysed using IGOR Pro (WaveMetrics, Version 6.3, Oregon, USA). A total of 180,050 dives were analysed over the 9-year period.

Each dive deeper than 1 m was identified, and different metrics were automatically calculated with a purpose-written macro in IGOR Pro. Among the parameters automatically extracted by the macro, we principally investigated dive depth and duration, the bottom phase duration, the number of undulations during the bottom phase in a dive and the post dive duration (see Ropert-Coudert, Wilson, Yoda, & Kato, 2007 for parameter definitions). Two indices were also calculated: diving efficiency (Ydenberg & Clark, 1989; Equation 1) and ACPUE (Attempts of Catch Per Unit Effort; Zimmer et al., 2010; Equation 2).

$$\text{Diving efficiency} = \frac{\text{Bottom duration}}{\text{Dive duration} + \text{Surface duration}}, \quad (1)$$



**FIGURE 1** Map showing the location of the Dumont d'Urville research station, along with maps of sea-ice concentration (SIC) for two different days within the same season 1995–1996, (a) 15th December 1995 and (b) 15th January 1996. White represents 100% sea ice covered and black represents open water. Black dot in (a) and white dot in (b) represent the studied penguin colony in Dumont D'Urville

$$\text{ACPUE} = \frac{\text{Number of undulations}}{\text{Bottom duration}} \quad (2)$$

Differences in sampling intervals (5 s in 1995 and 2007 and 1 s for all other years, Supporting Information Table S1), introduce a bias in the analysis of diving parameters. Therefore, data collected in 1995 and 2007 were excluded from most of the analyses presented, except when conducting the analysis of the maximum dive depth (considering deep dives only [ $>15$  m, see below]) and the fractal analysis on diving time series, as the method is robust to such changes in sampling interval (MacIntosh, Pelletier, Chiaradia, Kato, & Ropert-Coudert, 2013). Concerning the maximum dive depth, all 9 years were compared using only dives deeper than 15 m. This

threshold was extracted from the relationship between the maximum dive depth and the dive duration applied to all years and corresponds to a breakpoint dividing the distribution in two main portions (one concerning deep dives and one concerning surface dives). The technique assumes that these two categories of dives emerge from two different processes: shallow dives reflecting transit movements and deeper dives reflecting foraging movements (Ropert-Coudert et al., 2001). We have used the “segmented” package in R (Muggeo, 2015) to identify this threshold.

Student *t* tests were used to compare the means of each diving parameter when sample variances were equal (*F* test for equality of variances: *p*-value  $>0.05$ ) and Welch tests were used in case of non-

equal variances. The Bonferroni correction was applied to correct the level of significance because multiple comparisons were performed simultaneously.

Dive metrics were also modelled as a function of sea ice and bird ID, where SIC was modelled as a fixed effect and bird ID as a random effect. Linear Mixed Models (LMMs) combining fixed (SIC) and random (bird ID) effects were performed using the `lme` function of the R package “nlme” (Pinheiro, Bates, DebRoy, & Sarkar, 2016) and Generalized Linear Mixed Models (GLMMs) were applied using the `glmer` function of the R package “lme4” (Bates, Maechler, Bolker, & Walker, 2015). Dive duration and bottom duration were excluded from the analyses, as they were highly correlated to the dive depth and the number of undulations, respectively (with a Pearson correlation coefficient of 0.854 and 0.751, respectively). Diving efficiency and ACPUE (response variables) were analysed in a LMM with a normal error distribution in relation to SIC (fixed effect) and bird ID (random effect). To investigate the relationship between both maximum dive depth and post-dive duration (response variables) and sea ice using a LMM, the values of maximum dive depth and postdive duration were  $\log_{10}$ -transformed, while keeping SIC and bird ID as fixed and random predictors. The number of undulations was analysed as the response variable in a GLMM specified with a Poisson error distribution, with SIC as a fixed effect and bird ID as a random effect. The residuals from the chosen statistical models were visually inspected to ensure that the assumptions of residual homogeneity and homoscedasticity were met. For all diving metrics, we identified the best distribution to use (Normal,  $\log_{10}$ , Poisson, etc.) using the AIC. Then, a random intercept and slope model was performed assuming that the relationship between each diving metric and sea ice is different for each bird, and this model structure was always preferred compared to a simple intercept model (model selection still performed using the AIC). When a hump-shaped curve was observed, models specified with and without a quadratic term were compared (using the AIC) to investigate whether or not the relationship between the diving metric and SIC is statistically quadratic. For mixed models, the  $R^2$  can be divided in two components: the marginal  $R^2$  ( $R^2_m$ ) describes the proportion of variance explained by the fixed factor (SIC), while the conditional  $R^2$  ( $R^2_c$ ) describes the proportion of variance explained by both the fixed and random factors (SIC and bird ID). This permits to identify the part of the variance that is due to the fixed external effect (i.e. sea ice) and the part that is due to internal factors (i.e. the individual “quality”, characterised by the bird ID).

Penguins dive in foraging bouts, which correspond to sequences of multiple dives in succession over a certain period of time (Naito, Asaga, & Ohyama, 1990). Between two foraging bouts, individuals can rest at the surface, on land, on sea ice or transit to other foraging areas (Le Boeuf & Laws, 1994). Bout metrics are expected to reflect prey availability, as greater numbers of dives in a bout should coincide with larger prey patches (Boydlan, 1996; Sommerfeld et al., 2015). Similarly, small delays between dives within a bout likely reflect a higher prey patch density, while the distance between bouts (i.e. the distance between two prey patches) can be

linked to the prey encounter rate of the bird (Boydlan, 1996; Sommerfeld et al., 2015; Watanabe, Ito, & Takahashi, 2014). Here, foraging bouts were defined using the log survivorship analysis, which corresponds to a graphical method to specify the minimum surface interval between two dives belonging to separate bouts, also called the Bout Criterion Interval (BCI; Martin & Bateson, 1993). A single BCI value was calculated per year, based on the average BCI values of all birds in each year. We used the “segmented” package in R (Muggeo, 2015) to find BCIs (i.e. the “breakpoint” of the post-dive duration distribution). Following the definition of bouts, we calculated, for days with complete records only (24 hr), the number of dives per day, the number of dives per bout, the number of bouts per day, the bout duration and the mean bottom duration per bout for each bird. Mean comparisons between years were performed on the number of dives per day and the number of bouts per day using Student *t* tests and Welch tests applied with the Bonferroni correction. Because of the small amount of observations for each bird concerning these two parameters, the quadratic aspect of the relationship between both the number of dives and the number of bouts per day and SIC was tested comparing a GAM with and without a quadratic term. However, bout duration and bottom duration per bout were analysed using LMMs. Postdive duration within bouts (postdive duration < BCI) and postdive duration between bouts (interbout duration, postdive duration > BCI) were investigated separately. Mean comparisons between years were also performed on these parameters using Student *t* tests and Welch tests applied with the Bonferroni correction. To investigate the relationship between both postdive duration within bouts and interbout duration (response variables) and sea ice using a LMM, the values were  $\log_{10}$ -transformed, while keeping SIC and bird ID as fixed and random predictors. When a hump-shaped curve was observed, the chosen model specified with and without a quadratic term were compared (using the AIC) to investigate whether the relationship between the diving metric and SIC is statistically quadratic or not.

Finally, we also used fractal analysis (Mandelbrot, 1977) to investigate how the temporal organisation of dive sequences changes in relation to sea ice (see review in MacIntosh, 2014). Fractal time series analyses of animal behaviour aim to describe the structure of behaviour as it occurs through time, and to measure through a simple index the level of complexity observed in behavioural sequences (Asher et al., 2009; MacIntosh, 2014). Interactions between the behavioural strategies of an animal and the prevailing environmental conditions lead to the emergence of observed complexity signatures, which might reflect behavioural adaptations to environmental changes (Cribb & Seuront, 2016). Based on previous research (MacIntosh, Alados, & Huffman, 2011; Meyer et al., 2017; Sims et al., 2008), we hypothesized that foraging sequences would display greater complexity (here decreased long-range dependence and thus increased stochasticity) under more challenging and/or heterogeneous environmental conditions related to sea-ice cover. Following the method described by MacIntosh et al. (2013) and applied to Adélie penguins in Cottin et al. (2014) and Meyer, MacIntosh, Kato,

Chiaradia, and Ropert-Coudert (2015), we used Detrended Fluctuation Analysis (DFA; Peng et al., 1992) to measure long-range dependence in the sequential distribution of dives and surface times as an indicator of complexity in individual diving sequences. We performed DFA using the “fractal” package (Constantine & Percival, 2011) in R to estimate the scaling exponents ( $\alpha_{DFA}$ ) of these sequences (Peng et al., 1992), which measures the degree to which time series are long-range dependent and statistically self-similar (Taqqu, Teverovsky, & Willinger, 1995). This scaling exponent is also theoretically inversely related to the fractal dimension of a time series, which is a classical index of structural complexity (see Mandelbrot, 1977 for details). As  $\alpha_{DFA}$  increases from 0.5 to 1, diving sequences are increasingly persistent and long-range dependent, and the patterns as they occur over time are increasingly consistent/invariable (i.e. dives and surface times of a given length will be followed by dives and surface times of a similar length; Peng & Havlin, 1995). A GAM was performed to investigate variations in  $\alpha_{DFA}$  values (one value per individual) in relation to the overall SIC. The GAM was specified with a Gaussian family and five knots as there were more than 100 observations (121 values of  $\alpha_{DFA}$ ), following the optimal amount of smoothing recommended by Zuur et al. (2009). The GAM provides evidence of a quadratic relationship between  $\alpha_{DFA}$  and the overall SIC, GAMs with and without a quadratic term were compared using AIC.


The alpha level for all significance tests was set at 0.05. Results are generally presented as mean  $\pm$  standard errors (SE).

### 3 | RESULTS

#### 3.1 | Breeding success and sea ice

A significant effect of the overall SIC on breeding success was found by fitting a GAM specified with a quadratic relationship to the time series of breeding success ( $F$  test = 7.277, adjusted  $R^2 = 0.39$  and  $p$ -value = 0.015; Figure 2a). Breeding success was the highest in intermediate years, around 20% of SIC. The breeding season in 2014 had intermediate SIC values, but sea-ice retreat was so slow in that year that the distance between the colony and the open water was ca. 100 km in January. Birds had therefore to walk a very long way between feeding and nesting grounds, which likely explains the low breeding success in 2014. For this is reason, the GAM specified with a quadratic term was also performed excluding data from the year 2014, thereby improving the model and highlighting the quadratic relationship between breeding success and the overall SIC ( $F$  test = 16.78, adjusted  $R^2 = 0.578$  and  $p$ -value =  $8.43 \times 10^{-4}$ ; Figure 2b).

Daily SICs varied across years for the days when diving data were recorded (Figure 2). The years 1998 and 2001 were visually identified as low sea-ice cover years, while 2011 and 2012 corresponded to high sea-ice cover years (Figure 2). Other years are subsequently referred to as “intermediate” years. For the entire study period (1995–2014), both breeding success and SIC varied substantially (Figure 2).



**FIGURE 2** Fitted GAM results on Adélie penguins' breeding success using (a) all years (i.e. 20 years between 1995 and 2014) and (b) all years except 2014, in relation to the overall sea-ice concentration. Shades indicate 95% confidence intervals. Dark dots represent the studied years and light dots correspond to additional breeding success data concerning the period of interest 1995–2014

#### 3.2 | Foraging activity and sea ice


During the chick-guard stage, foraging trips typically last 1–2 days (mean  $\pm$  SE:  $49 \pm 51$  hr). The overall SIC had a significant effect in all selected models that investigated diving parameters as a function of sea ice (all  $p$ -values  $< 0.05$ ; Table 1). In all models, the random effect (due to individual variability) largely contributed to explain the global variability (Table 1).

With the exception of 2011 showing a high overall sea-ice concentration, the maximum dive depth was significantly greater, on average, during intermediate years (around 20% of sea-ice concentration) than during years with high or low sea-ice cover (model considering only deep dives from all years; Figure 3; Table 1). The shallowest average diving depths were observed in 2001 and 2012, with a mean of  $13.93 \pm 0.12$  and  $13.83 \pm 0.08$  m, respectively.

**TABLE 1** Results of the main mixed effects models for each diving metric in relation to the overall SIC

	Without $R^2c$	Quadratic $R^2m$	Effect $p$ -value	With $R^2c$	Quadratic $R^2m$	Effect $p$ -value	$\delta$ AIC
Dive depth (>15 m, 9 years)	0.46	0.06	0.0010	0.47	0.05	0.0010	2
Diving efficiency	0.17	0.02	<0.0001	0.15	0.03	<0.0001	18
Attempts of Catch Per Unit Effort	—	—	n.s.	—	—	n.s.	—
Number of undulations in a dive	—	—	n.s.	—	—	n.s.	—
Postdive duration within bouts	0.11	0.01	0.0301	0.10	0.01	0.0177	1
Interbout duration	0.13	0.01	0.0027	—	—	—	—


Note. The expression “n.s.” stands for “nonsignificant” and hyphens stands for “noncalculated”.  $\delta$ AIC represents the difference between AIC for the model without the quadratic effect and the one including it. For the maximum dive depth, models were performed on 49,411 deep dives (>15 m) of 121 birds over the 9-year period. For the other parameters, all 167,672 dives (>1 m) of 102 birds over the 7-year period (all years excluding 1995 and 2007) were analysed.



**FIGURE 3** Boxplots of dive depth according to years ordered by increasing overall sea-ice concentration (all years only considering dives deeper than 15 m). Medians and the 25th and 75th percentiles are represented by the box. The dots represent outliers. In each box, the central mark indicates the median, and the bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers, and the dots represent outliers that are plotted individually

Linear mixed models performed on all years with log-transformed maximum dive depth as the response variable and SIC as the predictor with and without the specification of a quadratic term were statistically similar based on the AIC criterion (AIC = 2,441 and 2,439, respectively; Table 1). Therefore, according to this model, intermediate and high SIC years are not statistically different. The same models were performed excluding the year 2011. In this case, the quadratic aspect of the relationship between the maximum depth and SIC was statistically evidenced. Indeed, the model with the quadratic effect was significantly better (AIC = 13,163,  $F$ -value = 17.192,  $R^2c$  = 0.288,  $R^2m$  = 0.042,  $p$ -value = 0.0001) than the one without it (AIC = 13,170,  $F$ -value = 9.931,  $R^2c$  = 0.287,  $R^2m$  = 0.026,  $p$ -value = 0.0021).


Diving efficiency followed a roughly open upward quadratic function, though the differences between years seemed small, with



**FIGURE 4** Boxplots of diving efficiency according to years ordered by increasing sea-ice concentration. Only the years with a sampling frequency of 1 s are considered here. In each box, the central mark indicates the median, and the bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers


lower diving efficiencies observed in intermediate years and higher efficiencies in high and low sea-ice cover years (Figure 4). The linear mixed model applied with a normal distribution performed with diving efficiency as the response variable and SIC as the predictor specified with a quadratic effect was significantly better than the one without it (AIC = -10,844 and -10,826, respectively; Table 1). Therefore, the model shows that a quadratic relationship exists between diving efficiency and the overall SIC ( $F$ -value = 25.966,  $R^2c$  = 0.15,  $R^2m$  = 0.03,  $p$ -value < 0.0001; Table 1).

We also tried to simply relate yearly mean diving efficiencies to yearly mean breeding success by fitting a linear model on these variables and found that the relationship was significantly linear ( $F$  = 8.97,  $adj$ - $R^2$  = 0.562,  $p$ -value = 0.02011). Indeed, these variables are highly negatively correlated (with a Pearson correlation coefficient of -0.749). In contrast, there was no significant effect of SIC on the number of undulations in a dive, ACPUE, mean bout duration and mean bottom duration per bout (Table 1).



**FIGURE 5** Barplots of (a) the mean number of dives per day and (b) the mean number of bouts per day, according to years ordered by increasing sea-ice concentration. Error bars represent standard errors. The asterisks refer to a statistical difference between two groups defined with brackets. The letters refer to groups that are not statistically different. Results are given for the 7 years with a sampling frequency of 1 s and considering all dives


Regarding the bout analysis, a total of 2,981 bouts were identified over the 7 years with a sampling frequency of 1 s, for which the BCI averaged 207.7 s (range: 168.2–247.5 s). Both the average numbers of dives and bouts per day were the highest in 2001 and 2012 ( $649.9 \pm 43.9$  and  $633.7 \pm 50.3$  dives per day and  $13.2 \pm 1.2$  and  $13.2 \pm 0.9$  bouts per day, for 2001 and 2012 respectively; Figure 5a,b). The lowest number of dives per day was found in 2014 ( $417.2 \pm 50.8$  dives per day) and the year 2010 showed the lowest number of bouts per day ( $7.0 \pm 0.8$  bouts per day; Figure 5a,b). GAMs were performed on all years using the number of dives per day as the response variable and the overall SIC as the predictor, with and without including a quadratic term. The model with the quadratic term was significantly better (AIC = 1,218,  $F$ -value = 4.537,  $R^2$ -adj = 0.079,  $p$ -value = 0.0360) than the model without it (AIC =



**FIGURE 6** Barplots representing (a) the post-dive duration within bouts (post-dive duration  $\leq 207$  s) and (b) the post-dive duration between bouts (post-dive duration  $> 207$  s, or interbout duration). Error bars represent standard errors. The asterisks refer to a statistical difference between two groups defined with brackets. The letters refer to groups that are not statistically different. Only 7 years with a sampling frequency of 1 s are considered here

1,224,  $F$ -value = 1.997,  $R^2$ -adj = 0.0505,  $p$ -value = 0.108) even though it did not explain much variation. Years with intermediate SIC were characterized by lower numbers of dives per day and lower numbers of bouts per day, on average (Student's  $t$  tests,  $p$ -values  $< 0.05$ ; Figure 5a,b). In contrast, for the number of bouts per day, the model without the quadratic term was significantly better (AIC = 521,  $F$ -value = 3.096,  $R^2$ -adj = 0.155,  $p$ -value = 0.0142) than the model including it (AIC = 531,  $F$ -value = 2.631,  $R^2$ -adj = 0.031,  $p$ -value = 0.108), thereby suggesting the absence of a quadratic effect for the overall SIC.

The model performed with post-dive duration within bouts (post-dive duration  $< 207$  s) as the response variable and SIC as the predictor, specified with a quadratic effect, was similar than the one without it (AIC = 30,603 and 30,604, respectively; Table 1), thereby



**FIGURE 7** Fitted GAM results of  $\alpha_{DFA}$  performed on (a) all years and (b) all years except 2001, considering all dives according to years classed by increasing sea-ice concentration

not highlighting the existence of a quadratic relationship between postdive duration within bouts and overall SIC ( $F$ -value = 7.622,  $R^2_c = 0.0916$ ,  $R^2_m = 0.008$ ,  $p$ -value = 0.0069; Figure 6a). Similarly, inter-bout durations appeared to be shorter in years of intermediate sea-ice conditions, on average (Welch test:  $p$ -value < 0.002), with the notable exception of 2014, compared with values obtained for high and low sea-ice cover years (Figure 6b), but no quadratic relationship could be evidenced for this parameter.

Concerning the fractal analysis, values of  $\alpha_{DFA}$  were similar to those reported previously for Adélie penguins at  $0.94 \pm 0.005$  (Meyer et al., 2015), indicating that foraging time series were long-range dependent and persistent ( $\alpha_{DFA} > 0.5$ , i.e. dive and post-dive times of a given length were more likely to be followed by dive and postdive times of similar lengths). The highest average value was recorded in 2011 ( $0.930 \pm 0.004$ ) and the lowest in 1998 ( $0.874 \pm 0.009$ ), with high and low SIC, respectively (Figure 7). Multiple  $t$  tests (applied with Bonferroni correction) comparing the year 2001 to other years have shown that the year 2001 significantly departed from this general trend ( $0.928 \pm 0.006$ ;  $t$  test:  $p$ -values < 0.001), considering the low SIC for this year. The  $\alpha_{DFA}$  tended to increase as SIC increased but with the year 2001 as an exception (Figure 7). A linear regression was performed using  $\alpha_{DFA}$  as the response variable and the overall SIC as the predictor. Results suggested that the relationship between  $\alpha_{DFA}$  and the overall SIC was nonlinear (linear model:  $F$  test = 6.4313,  $p$ -value = 0.0125,  $adj-R^2 = 0.0433$ ). Results depend significantly on whether or not we consider the year 2001 as an outlier. A GAM was performed on  $\alpha_{DFA}$  values according to the overall SIC for all years (Figure 7a). Results show that the effect of the overall SIC is significant ( $F$  test = 10.09,  $adj-R^2 = 0.254$ ,  $p$ -value =  $1.33 \times 10^{-6}$ ,  $AIC = -540.92$ ) but we did not observe a significant quadratic relationship (GAM with quadratic effect:  $F$  test = 7.076,  $adj-R^2 = 0.0561$ ,  $p$ -value = 0.0089,  $AIC = -518.13$ ). The same approach was applied to all years except 2001 (Figure 7b). The effect of the overall SIC remained significant

( $F$  test = 17.04,  $adj-R^2 = 0.341$ ,  $p$ -value =  $1.26 \times 10^{-8}$ ,  $AIC = -459.42$ ) but again, we did not observe a significant quadratic relationship (GAM with quadratic effect:  $F$  test = 16.68,  $adj-R^2 = 0.322$ ,  $p$ -value =  $9.08 \times 10^{-5}$ ,  $AIC = -457.49$ ).

In other words, because no quadratic trend could be evidenced in either scenario, it can only generally be assumed that the temporal organisation of diving behaviour is more consistent/less stochastic in the diving sequences (i.e. characterized by a higher degree of dependence on history) during high sea-ice cover years. Conversely, reduced sea-ice concentration coincided with more complex/more stochastic diving sequences.

## 4 | DISCUSSION

### 4.1 | Similar effect of sea-ice cover on breeding success and diving behaviour

The breeding success and some diving parameters of Adélie penguins at Dumont d'Urville varied quadratically with sea-ice cover, both reaching a peak – either upward or downward – when sea-ice cover is intermediate, around 20%. Our results confirm the existence of an “optimal” range of sea-ice cover for Adélie penguins, as suggested by Ainley (2002) and corroborated by Barbraud et al. (2015) for breeding success. Ballard et al. (2010) also found a quadratic effect of SIC on foraging parameters in Adélie penguins, showing an optimum SIC around 10%–15%. The difference from our findings of an optimum around 20% is noteworthy and probably results from the fact that these authors studied trip duration and total food delivered, while we investigated the influence of SIC on diving activity. In addition, the similarity in the shape of the curve and the coincidence of the peaks around a given range suggest that sea ice may influence breeding success via prey availability (prey abundance and accessibility), which might affect diving behaviour. During the years of intermediate sea-ice cover, breeding success was generally maximized,

suggesting that these years offer enhanced food availability compared to years with high or low sea-ice cover. Indeed, as krill feed on under-ice communities (e.g. microalgae; Nicol, 2006), low sea-ice cover would reduce the amount of food available to krill, therefore negatively affecting their abundance. Conversely, foraging costs for chick-rearing penguins increase when sea-ice cover is high, forcing parents to walk longer distances on ice to reach open foraging areas (Ropert-Coudert et al., 2015). This, in turn, increases body mass loss for parents and, as their foraging trips are longer, decreases the provisioning frequency to the chicks (Ballard et al., 2010; Davis, 1982). A recent study has also shown that in low sea-ice cover conditions, the visual performance of predators is enhanced by increased light (Langbehn & Varpe, 2017). Consequently, intermediate sea-ice conditions seem to maximize prey availability and to minimize foraging costs for penguins, thereby resulting in enhanced foraging success. The 20% sea-ice concentration could correspond to a Marginal Ice Zone (MIZ), which is recognized as a concentration enhancing productivity in the trophic chains (Smith, Bird, & Budgell, 1988; Smith & Nelson, 1985). However, our optimal SIC value is a mean calculated over the whole potential foraging area of the birds and over a season, and it would be surprising if this whole area was covered with MIZ all summer long.

Although interindividual variability was high and, consequently, the quadratic trend in diving parameters was not as clear as it was for breeding success, the same pattern was found in some of the diving parameters investigated. More importantly, this trend was evident at various temporal scales: at the scale of the foraging trip and at the finer scale of individual dives. During years of intermediate sea-ice conditions (characterized by high breeding success and thus supposedly enhanced prey availability), Adélie penguins showed lower diving efficiencies but dove deeper than in years with high or low sea-ice cover; and at a lesser extent, spent less time commuting between prey patches while performing fewer bouts per day and fewer dives per day. In other words, although located at deeper depths, these metrics strongly suggest that intermediate years are characterized by higher prey encounter rates and prey availability.

The decreasing complexity in foraging behaviour as sea-ice cover increases indicates that Adélie penguins become increasingly constrained (i.e. they do not have much flexibility) in their diving sequences with increasing sea-ice cover. Following Reynolds, Ropert-Coudert, Kato, Chiaradia, and Macintosh (2015), we can postulate that birds in heavy ice would favour exploitation over exploration, while it would be the reverse when water becomes increasingly free of ice. This finding is also in accord with the fact that in intermediate sea-ice cover years, Adélie penguins exhibited high levels of plasticity in foraging efficiency (Lescroël et al., 2014). Indeed, a greater heterogeneity in the vertical distribution of prey like that observed in open water (Pelletier, Kato, Chiaradia, & Ropert-Coudert, 2012; Ropert-Coudert, Kato, & Chiaradia, 2009) should force penguins to favour the exploration of their environment, which is more likely to lead to complex behaviour (Kembro, Perillo, Pury, Satterlee, & Marin, 2009; MacIntosh et al., 2011; Shimada, Minesaki, & Hara, 1995). In contrast, the consistent/invariable diving sequences occurring during

years with high sea-ice cover could be due to the fact that birds are more constrained in their foraging movements, targeting shallower prey patches for which the need for more exploratory dives is limited. The presence of polynyas under these conditions becomes crucial in explaining the foraging activity of penguins as the high predictability of prey in polynyas likely leads to more consistent/invariable diving sequences (Meyer, 2016; Meyer et al., 2017).

Interestingly, some years clearly deviated from the norm regarding the relationship between sea-ice concentration and breeding success (2014) and the relationship between sea-ice concentration and diving parameters (depth in 2011 and 2014, postdive duration in 2001, interbout duration in 2014, daily number of dives in 2011, daily number of bouts in 2011). As 2001 and 2011 were characterized by the lowest and highest sea-ice concentrations (14% and 32%, respectively) in our study, it is possible that the effects of sea-ice concentration on certain characteristics of the diving behaviour of penguins are reversed over a certain threshold. However, 2014 was not characterized by such extreme sea-ice concentrations; this year showed a late sea-ice retreat and a distance between the colony and the open water around 100 km in January. This suggests that these environmental factors are able to disrupt the effects of sea-ice concentration on the foraging behaviour and the breeding performance of penguins. However, some diving parameters (diving efficiency,  $\alpha$ DFA) were not affected by these environmental factors, and therefore better reflected sea-ice changes. To better anticipate how changes in sea-ice concentration will affect Adélie penguins in the coming decades, future studies should examine how it interacts with other environmental changes to modulate their foraging behaviour and breeding success.

## 4.2 | The paradox of the diving efficiency index

If years with intermediate sea-ice cover correspond to profitable prey availability and higher breeding success, why was diving efficiency significantly lower during these years? This is indeed surprising as breeding success has been linked to foraging success (meal size provided to the chicks and their fledging mass) in Adélie penguins (Clarke, Knowles, Irvine, & Phillips, 2002). The explanation probably lies in the parameters involved in the calculation of this index. Besides the fact that undulations in the dive profiles may not always reflect prey capture (Watanabe & Takahashi, 2013; Watanabe et al., 2014), the index was initially designed for shallow, benthic feeders, for which prey items are predictably located at the bottom of the water column. Yet, the term “efficiency” has often been applied to species for which prey locations are not predictable in the water column (as in our case, but see also Zimmer et al., 2010), making variation in diving “efficiency” difficult to interpret. In fact, this index informs us about how birds deal with the different components of a dive (transit, hunting and resting) within a limited time window. Furthermore, both the quality and the quantity of the encountered prey are necessary to understand the mechanisms underlying the different components of a dive. During periods of high sea-ice concentration, the shallow diving activity of penguins suggests that they search for

prey just under the ice. Do penguins target young krill or are they switching to fish associated with sea ice? The question deserves further investigation. In contrast, in a year of low sea-ice cover, we hypothesize that prey might be less available, as their area of occurrence (the sea-ice edge) is less extensive and prey resources could be depleted quickly as birds concentrate on the ice edges. From the above, we can hypothesize that sea ice at moderate amounts is favourable to the prey itself and this optimal is reflected in the activity of the penguins. Adélie penguins probably forage on highly profitable albeit deeper prey fields during years of intermediate sea-ice conditions, which translates in an apparent lower diving efficiency during those years due to the metrics used in its calculation. Regardless of the mechanisms involved, we strongly advocate for the diving efficiency index to be renamed so as to better reflect the heterogeneous nature of the phenomenon it measures.

### 4.3 | Perspectives

Our results suggest the existence of an optimal range of sea-ice cover that enhances the foraging success and subsequently the breeding success of Adélie penguins, peaking at approximately 20% cover in the foraging zone around DDU (Ainley, 2002; see also Barbraud et al., 2015). There may be other factors that can impact penguin breeding success. Optimal sea-ice conditions are likely to be a necessary but insufficient condition for high breeding success (Youngflesh et al., 2017). For instance, the timing of phytoplankton blooms may influence the availability and quality of prey (Atkinson et al., 2008; Saba et al., 2014). In addition, climatic conditions such as low ambient air temperature or high levels of precipitation can also affect the survival of the chicks (Boersma & Rebstock, 2014; Ropert-Coudert et al., 2015), directly impacting breeding success. Delimiting precisely the extent of this optimal range will require further investigation but is an essential step in assessing the capacity of this species to buffer forthcoming changes in sea-ice conditions. One can hypothesize that the narrower the range of optimal sea-ice conditions, the more affected the eco-indicating species will be by environmental changes and other potential threats, such as marine pollution, expanding tourism or other anthropogenic activities, the increasing risk of competing with fisheries or introduced species exploiting krill resources (Kennicutt et al., 2014; Xavier et al., 2016). To better refine the limits of this range, including additional years in the monitoring effort is a prerequisite, but homogenising the protocol of data collection is also paramount. It would also be interesting to add a horizontal dimension (e.g. GPS tracking, see Widmann et al., 2015) to the monitoring of the foraging behaviour of penguins, to precisely assess the ice habitats used by penguins (polynyas, fast ice cracks, etc.).

Finally, it seems necessary to extend this approach to other colonies of Adélie penguins around Antarctica because changes in sea-ice conditions in some parts of Western Antarctica are opposite to what is currently being observed at Dumont d'Urville (e.g. Antarctic Peninsula, Lima & Estay, 2013; McClintock, Ducklow, & Fraser, 2008; Stammerjohn, Massom, Rind, & Martinson, 2012). If the situation observed in the Peninsula – less sea ice links to population decrease – becomes

a general trend around the continent, the global Adélie penguin populations might soon be facing unprecedented challenges. Yet, penguins may be capable of greater flexibility than current knowledge suggests. Results of paleoecological studies show that Adélie penguins have not always been the creatures of the sea ice that they are considered to be today. Indeed, they have switched from eating fish to krill after humans began to hunt whales in the past (Emslie & Patterson, 2007). By removing krill predators from the environment, the penguins took advantage of an open niche to target and exploit krill surpluses. Such flexibility indicates the complexity of investigating trophic links in the so-called simple Southern Ocean ecosystem. In the framework of the CCAMLR Ecosystem Monitoring Program (CCAMLR, 2014), this calls for the need to use collaborations among existing Adélie penguin monitoring sites to determine in real time the changes occurring in these contrasted environments (using tracking data among other variables), and to better evaluate the range of impacts these changes have on ecosystem functioning. At a time when the CCAMLR is considering the feasibility of developing a network of Marine Protected Areas around the continent, such monitoring sites would be particularly useful to test the efficiency of the MPA in protecting the resources and the species that depend on them.

### ACKNOWLEDGEMENTS

The data used in this work were collected by members of various scientific programs supported by the French Polar Institute Paul-Emile Victor: programmes 137 administered by Dr. Y. Le Maho (diving data from 1995, 1998, 2007, 2009), 1,091 administered by Dr. Y. Ropert-Coudert (diving data from 2010 to 2015), and 109 administered by Dr. H. Weimerskirch (breeding success data from 1995 to 2015). We thank all the volunteers in the field, who participated to the sampling effort and/or the data processing of the data, especially A. Ancel, J. Baudat, K. Yoda, M. Cottin, A. M. Thierry, M. Spée, T. Poupart, M. Pellé, B. Arthur, F. Angelier, O. Chastel, F. Amélineau, M. Debin, W. Bonneau, D. Rodary, and all the members of the logistic on site for their assistance. We also thank Dr. J. Hinke and three anonymous referees for their helpful comments on the manuscript. This study was financially supported by the following institutions: the WWF-UK through R. Downie, the Japanese Mombukagakusho and the Japanese Society for the Promotion of Science, the Zone Atelier Antarctique et Subantarctique –LTER France of the CNRS. This study was approved by the ethic committee of the French Polar Institute Paul Emile Victor.

### ORCID

Camille Le Guen  <http://orcid.org/0000-0002-2365-5283>

### REFERENCES

- Ainley, D. G. (2002). *The Adélie Penguin: Bellwether of climate change* (310 pp.). New York, NY: Columbia University Press. <https://doi.org/10.7312/ainl12306>

- Ainley, D. G., Wilson, P. R., Barton, K. J., Ballard, G., Nur, N., & Karl, B. (1998). Diet and foraging effort of Adélie penguins in relation to pack-ice conditions in the southern Ross Sea. *Polar Biology*, 20, 311–319. <https://doi.org/10.1007/s003000050308>
- Asher, L., Collins, L. M., Ortiz-Pelaez, A., Drewe, J. A., Nicol, C. J., & Pfeiffer, D. U. (2009). Recent advances in the analysis of behavioural organization and interpretation as indicators of animal welfare. *Journal of the Royal Society, Interface/the Royal Society*, 6, 1103–1119. <https://doi.org/10.1098/rsif.2009.0221>
- Atkinson, A., Siegel, V., Pakhomov, E. A., Rothery, P., Loeb, V., Ross, R. M., ... Fleming, A. H. (2008). Oceanic circumpolar habitats of Antarctic krill. *Marine Ecology Progress Series*, 362, 1–23. <https://doi.org/10.3354/meps07498>
- Ballard, G., Dugger, K., Nur, N., & Ainley, D. (2010). Foraging strategies of Adélie penguins: adjusting body condition to cope with environmental variability. *Marine Ecology Progress Series*, 405, 287–302. <https://doi.org/10.3354/meps08514>
- Barbraud, C., Delord, K., & Weimerskirch, H. (2015). Extreme ecological response of a seabird community to unprecedented sea ice cover. *Royal Society Open Science*, 2, 140456. <https://doi.org/10.1098/rsos.140456>
- Barbraud, C., & Weimerskirch, H. (2003). Climate and density shape population dynamics of a marine top predator. *Proceedings of the Royal Society B: Biological Sciences*, 270, 2111–2116. <https://doi.org/10.1098/rspb.2003.2488>
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1–48.
- Beaulieu, M., Dervaux, A., Thierry, A.-M., Lazin, D., Le Maho, Y., Ropert-Coudert, Y., ... Ancel, A. (2010). When sea-ice clock is ahead of Adélie penguins' clock. *Functional Ecology*, 24, 93–102. <https://doi.org/10.1111/j.1365-2435.2009.01638.x>
- Boersma, P., & Rebstock, G. (2014). Climate change increases reproductive failure in Magellanic Penguins. *PLoS One*, 9, e85602. <https://doi.org/10.1371/journal.pone.0085602>
- Bost, C. A., Cott, C. É., Bailleul, F., Cherel, Y., Charrassin, J. B., Guinet, C., ... Weimerskirch, H. (2009). The importance of oceanographic fronts to marine birds and mammals of the southern oceans. *Journal of Marine Systems*, 78, 363–376. <https://doi.org/10.1016/j.jmarsys.2008.11.022>
- Boyd, I. L., & Murray, A. W. A. (2001). Monitoring a marine ecosystem using upper trophic level predators. *Journal of Animal Ecology*, 70, 747–760. <https://doi.org/10.1046/j.0021-8790.2001.00534.x>
- Boydlan, L. (1996). Temporal scales of foraging in a marine predator. *Ecology*, 77, 426–434.
- Brierley, A. (2002). Antarctic krill under sea ice: elevated abundance in a narrow band just south of ice edge. *Science*, 295, 1890–1892. <https://doi.org/10.1126/science.1068574>
- Cavaliere, D., Parkinson, C., Gloersen, P., & Zwally, H. (1996, updated yearly). *Sea ice concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS passive microwave data*. Boulder, CO: NASA National Snow and Ice Data Center Distributed Active Archive Center. <https://doi.org/10.5067/8gq8lqzvl0vl>
- CCAMLR (2014). *CCAMLR ecosystem monitoring program standard methods*. Hobart, TAS: CCAMLR.
- Cherel, Y. (2008). Isotopic niches of emperor and Adélie penguins in Adélie Land, Antarctica. *Marine Biology*, 154, 813–821. <https://doi.org/10.1007/s00227-008-0974-3>
- Clarke, A., & Harris, C. M. (2003). Polar marine ecosystems: major threats and future change. *Environmental Conservation*, 30, 1–25.
- Clarke, J., Knowles, K., Irvine, L., & Phillips, B. (2002). Chick provisioning and breeding success of Adélie penguins at Béchervaise Island over eight successive seasons. *Polar Biology*, 25, 21–30. <https://doi.org/10.1007/s003000100307>
- Constantine, W., & Percival, D. (2011). Fractal: fractal time series modeling and analysis. R package v. 1.1-1.
- Cottin, M., MacIntosh, A., Kato, A., Takashi, A., Debin, M., Raclot, T., & Ropert-Coudert, Y. (2014). Corticosterone administration leads to a transient alteration of foraging behaviour and complexity in a diving seabird. *Marine Ecology Progress Series*, 496, 249–262. <https://doi.org/10.3354/meps10618>
- Cottin, M., Raymond, B., Kato, A., Amélineau, F., Le Maho, Y., Raclot, T., ... Ropert-Coudert, Y. (2012). Foraging strategies of male Adélie penguins during their first incubation trip in relation to environmental conditions. *Marine Biology*, 159, 1843–1852. <https://doi.org/10.1007/s00227-012-1974-x>
- Cribb, N., & Seuront, L. (2016). Changes in the behavioural complexity of bottlenose dolphins along a gradient of anthropogenically-impacted environments in South Australian coastal waters: Implications for conservation and management strategies. *Journal of Experimental Marine Biology and Ecology*, 482, 118–127. <https://doi.org/10.1016/j.jembe.2016.03.020>
- Croxall, J., & Lishman, G. S. (1987). The food and feeding ecology of penguins. In J. Croxall (Ed.), *Seabirds: Feeding ecology and role in marine ecosystems* (pp. 101–133). Cambridge, UK: Cambridge University Press.
- Croxall, J. P., Trathan, P. N., & Murphy, E. J. (2002). Environmental change and Antarctic seabird populations. *Science*, 297, 1510–1514. <https://doi.org/10.1126/science.1071987>
- Davis, L. (1982). Timing of nest relief and its effect on breeding success in Adélie penguins (*Pygoscelis adeliae*). *Condor*, 84, 178–183. <https://doi.org/10.2307/1367665>
- Doney, S. C., Ruckelshaus, M., Duffy, J. E., Barry, J. P., Chan, F., English, C. A., ... Talley, L. D. (2012). Climate change impacts on marine ecosystems. *Annual Review of Marine Science*, 4, 11–37. <https://doi.org/10.1146/annurev-marine-041911-111611>
- Dugger, K. M., Ballard, G., Ainley, D. G., Lyver, P. O. B., & Schine, C. (2014). Adélie penguins coping with environmental change: results from a natural experiment at the edge of their breeding range. *Frontiers in Ecology and Evolution*, 2, 1–12.
- Elton, C. (1958). *The ecology of invasions by animals and plants* (196 pp.). London, UK: Methuen edn. <https://doi.org/10.1007/978-1-4899-7214-9>
- Emmerson, L., & Southwell, C. (2008). Sea ice cover and its influence on Adélie penguin reproductive performance. *Ecology*, 89, 2096–2102. <https://doi.org/10.1890/08-0011.1>
- Emslie, S., & Patterson, W. P. (2007). Abrupt recent shift in  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in Adélie penguin eggshell in Antarctica. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 11666–11669. <https://doi.org/10.1073/pnas.0608477104>
- Fraser, W., Trivelpiece, W., Ainley, D. G., & Trivelpiece, S. (1992). Increases in Antarctic penguin populations: reduced competition with whales or a loss of sea ice due to environmental warming? *Polar Biology*, 11, 525–531.
- Frederiksen, M., Mavor, R. A., & Wanless, S. (2007). Seabirds as environmental indicators: The advantages of combining data sets. *Marine Ecology Progress Series*, 352, 205–211. <https://doi.org/10.3354/meps07071>
- Furness, R. W., & Camphuysen, C. J. (1997). Seabirds as monitors of the marine environment. *ICES Journal of Marine Science*, 54, 726–737. <https://doi.org/10.1006/jmsc.1997.0243>
- Hijmans, R. J., van Etten, J., Cheng, J., Mattiuzzi, M., Sumner, M., Greenberg, J. A., ... Shortridge, A. (2016). Package “raster”. CRAN-R.2.5-8.
- Hindell, M., Bradshaw, C., Harcourt, R., & Guinet, C. (2003). Ecosystem monitoring: Are seals a potential tool for monitoring change in marine systems? In N. J. Gales & M. A. K. R. Hindell (Eds.), *Marine mammals. Fisheries, tourism and management issues* (pp. 330–343). Melbourne, Vic.: CSIRO Publishing.
- IPCC (2007). *Climate Change 2007 Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (104 pp.). Core Writing Team, R.K. Pachauri and A. Reisinger. Geneva, Switzerland: IPCC.

- Jennings, S., & Kaiser, M. (1998). The effects of fishing on marine ecosystems. *Advances in Marine Biology*, 34, 201–351. [https://doi.org/10.1016/S0065-2881\(08\)60212-6](https://doi.org/10.1016/S0065-2881(08)60212-6)
- Jenouvrier, S., Barbraud, C., & Weimerskirch, H. (2006). Sea ice affects the population dynamics of Adélie penguins in Terre Adélie. *Polar Biology*, 29, 413–423. <https://doi.org/10.1007/s00300-005-0073-6>
- Jenouvrier, S., Holland, M., Stroeve, J., Serreze, M., Barbraud, C., Weimerskirch, H., & Caswell, H. (2014). Projected continent-wide declines of the emperor penguin under climate change. *Nature Climate Change*, 4, 715–718. <https://doi.org/10.1038/nclimate2280>
- Kato, A., Ropert-Coudert, Y., & Naito, Y. (2002). Changes in Adélie penguin breeding populations in Lutzow-Holm Bay, Antarctica, in relation to sea-ice conditions. *Polar Biology*, 25, 934–938.
- Kato, A., Watanuki, Y., & Naito, Y. (2003). Annual and seasonal changes in foraging site and diving behavior in Adélie penguins. *Polar Biology*, 26, 389–395.
- Katsanevakis, S., Wallentinus, I., Zenetos, A., Leppäkoski, E., Çınar, M. E., Öztürk, B., ... Cardoso, A. C. (2014). Impacts of invasive alien marine species on ecosystem services and biodiversity: a pan-European review. *Aquatic Invasions*, 9, 391–423. <https://doi.org/10.3391/ai>
- Kembro, J., Perillo, M., Pury, P., Satterlee, D., & Marin, R. (2009). Fractal analysis of the ambulation pattern of Japanese quail. *British Poultry Science*, 50, 161–170. <https://doi.org/10.1080/00071660802710116>
- Kennicutt, M. C. 2nd, Chown, S. L., Cassano, J. J., Liggett, D., Massom, R., Peck, L. S., ... Sutherland, W. J. (2014). Six priorities for Antarctic science. *Nature*, 512, 23–25. <https://doi.org/10.1038/512023a>
- Knox, G. A. (2006). *Biology of the Southern Ocean* (2nd ed., 640 pp). Boca Raton, FL: CRC Press. <https://doi.org/10.1201/CRCMARINEBIO>
- Langbehn, T., & Varpe, Ø. (2017). Sea-ice loss boosts visual search: Fish foraging and changing pelagic interactions in polar oceans. *Global Change Biology*, 23, 5318–5330. <https://doi.org/10.1111/gcb.13797>
- Le Boeuf, B. J., & Laws, R. M. (1994). *Elephant seals: Population ecology, behavior and physiology* (414 pp.). Berkeley, CA: University of California Press.
- Lescroël, A., Ballard, G., Gremillet, D., Authier, M., & Ainley, D. G. (2014). Antarctic climate change: extreme events disrupt plastic phenotypic response in Adélie Penguins. *PLoS One*, 9, e85291. <https://doi.org/10.1371/journal.pone.0085291>
- Lescroël, A., Dugger, K. M., Ballard, G., & Ainley, D. G. (2009). Effects of individual quality, reproductive success and environmental variability on survival of a long-lived seabird. *Journal of Animal Ecology*, 78, 798–806. <https://doi.org/10.1111/j.1365-2656.2009.01542.x>
- Libertelli, M. M., Coria, N., & Marateo, G. (2003). Diet of the Adélie penguin during three consecutive chick rearing periods at Laurie Island. *Polish Polar Research*, 24, 133–142.
- Lima, M., & Estay, S. (2013). Warming effects in the western Antarctic Peninsula ecosystem: the role of population dynamic models for explaining and predicting penguin trends. *Population Ecology*, 55, 557–565. <https://doi.org/10.1007/s10144-013-0386-1>
- MacIntosh, A. J. J. (2014). The fractal primate: Interdisciplinary science and the math behind the monkey. *Primate Research*, 30, 95–119. <https://doi.org/10.2354/psj.30.011>
- MacIntosh, A. J., Alados, C., & Huffman, M. (2011). Fractal analysis of behaviour in a wild primate: behavioural complexity in health and disease. *Journal of the Royal Society, Interface / the Royal Society*, 8, 1497–1509. <https://doi.org/10.1098/rsif.2011.0049>
- MacIntosh, A. J. J., Pelletier, L., Chiaradia, A., Kato, A., & Ropert-Coudert, Y. (2013). Temporal fractals in seabird foraging behaviour: diving through the scales of time. *Scientific Reports*, 3, 1884. <https://doi.org/10.1038/srep01884>
- Mandelbrot, B. (1977). *The fractal geometry of nature*. New-York, NY: W.H. Freeman and Company.
- Martin, P., & Bateson, P. (1993). *Measuring behaviour: An introductory guide* (222 pp). Cambridge, UK: Cambridge University Press. <https://doi.org/10.1017/CBO9781139168342>
- McClintock, J. B., Ducklow, H. W., & Fraser, W. (2008). Ecological responses to climate change on the Antarctic Peninsula. *American Scientist*, 96, 302–310. <https://doi.org/10.1511/2008.73.3844>
- Meyer, X. (2016). Does complexity in behavioral organization allow seabirds to adapt to changes in their environment? (155 pp.). PhD Université de Strasbourg.
- Meyer, X., Macintosh, A. J. J., Chiaradia, A., Kato, A., Mattern, T., Sueur, C., & Ropert, Y. (2017). Shallow divers, deep waters and the rise of behavioural stochasticity. *Marine Biology*, 164, 149. <https://doi.org/10.1007/s00227-017-3177-y>
- Meyer, X., MacIntosh, A. J. J., Kato, A., Chiaradia, A., & Ropert-Coudert, Y. (2015). Hydrodynamic handicaps and organizational complexity in the foraging behavior of two free-ranging penguin species. *Animal Biotelemetry*, 3, 25. <https://doi.org/10.1186/s40317-015-0061-8>
- Mugge, V. M. R. (2015). Title Regression Models with Breakpoints/Changepoints Estimation. CRAN-R0.2-9.3.
- Naito, Y., Asaga, T., & Ohyama, Y. (1990). Diving behavior of Adélie penguins determined by Time-Depth Recorder. *The Condor*, 92, 582–586. <https://doi.org/10.2307/1368676>
- Nicol, S. (2006). Krill, currents, and sea ice: *Euphausia superba* and its changing environment. *BioScience*, 56, 111. [https://doi.org/10.1641/0006-3568\(2006\)056\[0111:KCASIE\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)056[0111:KCASIE]2.0.CO;2)
- Orians, G., & Pearson, N. (1979). On the theory of central place foraging. In D. Horn, R. Mitchell, & G. Stairs (Eds.), *Analysis of ecological systems* (pp. 154–177). Columbus, OH: The Ohio State University Press.
- Pelletier, L., Kato, A., Chiaradia, A., & Ropert-Coudert, Y. (2012). Can thermoclines be a cue to prey distribution for marine top predators? a case study with little penguins. *PLoS One*, 7, 4–8.
- Peng, C. K., Buldyrev, S. V., Goldberger, A. L., Havlin, S., Sciortino, F., Simons, M., & Stanley, H. E. (1992). Long-range correlations in nucleotide sequences. *Nature*, 356, 168–170. <https://doi.org/10.1038/356168a0>
- Peng, C., & Havlin, S. (1995). Quantification of scaling exponents and crossover phenomena in nonstationary heartbeat time series. *Chaos*, 5, 82–87. <https://doi.org/10.1063/1.166141>
- Piatt, J. F., Sydeman, W. J., & Wiese, F. (2007). Introduction: A modern role for seabirds as indicators. *Marine Ecology Progress Series*, 352, 199–204. <https://doi.org/10.3354/meps07070>
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team (2016). nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-128.
- R Development Core Team. (2015). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Ratcliffe, N., Takahashi, A., Oulton, C., Fukuda, M., Fry, B., Crofts, S., ... Trathan, P. N. (2014). A leg-band for mounting geolocator tags on penguins. *Marine Ornithology*, 42, 23–26.
- Reynolds, A., Ropert-Coudert, Y., Kato, A., Chiaradia, A., & Macintosh, A. J. J. (2015). A priority-based queuing process explanation for scale-free foraging behaviours. *Animal Behaviour*, 108, 67–71. <https://doi.org/10.1016/j.anbehav.2015.07.022>
- Richardson, A., & Polocsanska, E. (2008). Under-resourced, under threat. *Science*, 320, 1294–1295. <https://doi.org/10.1126/science.1156129>
- Ropert-Coudert, Y., Kato, A., Baudat, J., Bost, C.-A., Le Maho, Y., & Naito, Y. (2001). Feeding strategies of free-ranging Adélie penguins *Pygoscelis adeliae* analysed by multiple data recording. *Polar Biology*, 24, 460–466. <https://doi.org/10.1007/s003000100234>
- Ropert-Coudert, Y., Kato, A., & Chiaradia, A. (2009). Impact of small-scale environmental perturbations on local marine food resources: a case study of a predator, the little penguin. *Proceedings Biological Sciences/The Royal Society*, 276, 4105–4109. <https://doi.org/10.1098/rspb.2009.1399>
- Ropert-Coudert, Y., Kato, A., Gremillet, D., & Grenner, F. (2012). Bio-logging: recording the ecophysiology and behaviour of animals moving freely in their environment. In J.-F. Le Galliard, J.-M. Guarini & F. Gaill (Eds.), *Sensors for ecology: towards an integrated knowledge of ecosystems* (pp. 17–41). Paris, France: CNRS.

- Ropert-Coudert, Y., Kato, A., Meyer, X., Pellé, M., MacIntosh, A. J. J., Angelier, F., ... Raclot, T. (2015). A complete breeding failure in an Adélie penguin colony correlates with unusual and extreme environmental events. *Ecography*, *38*, 111–113. <https://doi.org/10.1111/ecog.01182>
- Ropert-Coudert, Y., & Wilson, R. (2005). Trends and perspectives in animal-attached remote sensing. *Frontiers in Ecology and the Environment*, *3*, 437–444. [https://doi.org/10.1890/1540-9295\(2005\)003\[0437:TAPIAR\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2005)003[0437:TAPIAR]2.0.CO;2)
- Ropert-Coudert, Y., Wilson, R. P., Yoda, K., & Kato, A. (2007). Assessing performance constraints in penguins with externally-attached devices. *Marine Ecology Progress Series*, *333*, 281–289. <https://doi.org/10.3354/meps333281>
- Saba, G. K., Fraser, W. R., Saba, V. S., Iannuzzi, R. A., Coleman, K. E., Doney, S. C., ... Schofield, O. M. (2014). Winter and spring controls on the summer food web of the coastal West Antarctic Peninsula. *Nature Communications*, *5*, 4318. <https://doi.org/10.1038/ncomms5318>
- Sedwick, P. N., & DiTullio, G. R. (1997). Regulation of algal blooms in Antarctic shelf waters by the release of iron from melting sea ice. *Geophysical Research Letters*, *24*, 2515–2518. <https://doi.org/10.1029/97GL02596>
- Shimada, I., Minesaki, Y., & Hara, H. (1995). Temporal fractal in the feeding behavior of *Drosophila melanogaster*. *Journal of Ethology*, *13*, 153–158. <https://doi.org/10.1007/BF02350106>
- Sims, D. W., Southall, E. J., Humphries, N. E., Hays, G. C., Bradshaw, C. J. A., Pitchford, J. W., ... Metcalfe, J. D. (2008). Scaling laws of marine predator search behaviour. *Nature*, *451*, 1098–1102. <https://doi.org/10.1038/nature06518>
- Smith, R. C., Ainley, D., Baker, K., Domack, E., Emslie, S., Fraser, B., ... Vernet, M. (1999). Marine ecosystem sensitivity to climate change. *BioScience*, *49*, 393–404. <https://doi.org/10.2307/1313632>
- Smith, D. IV, Bird, A., & Budgell, W. (1988). A numerical study of mesoscale eddy interaction with a marginal ice zone. *Journal of Geophysical Research*, *93*, 12461–12473. <https://doi.org/10.1029/JC093iC10p12461>
- Smith, W., & Nelson, D. (1985). Phytoplankton bloom produced by a receding ice edge in the Ross Sea: Spatial coherence with the density field. *Science*, *227*, 163–166. <https://doi.org/10.1126/science.227.4683.163>
- Sommerfeld, J., Kato, A., Ropert-Coudert, Y., Garthe, S., Wilcox, C., & Hindell, M. A. (2015). Flexible foraging behaviour in a marine predator, the Masked booby (*Sula dactylatra*), according to foraging locations and environmental conditions. *Journal of Experimental Marine Biology and Ecology*, *463*, 79–86. <https://doi.org/10.1016/j.jembe.2014.11.005>
- Stammerjohn, S., Massom, R., Rind, D., & Martinson, D. (2012). Regions of rapid sea ice change: An inter-hemispheric seasonal comparison. *Geophysical Research Letters*, *39*, L06501.
- Taqqu, M., Teverovsky, V., & Willinger, W. (1995). Estimators for long-range dependence: an empirical study. *Fractals*, *3*, 785–788. <https://doi.org/10.1142/S0218348X95000692>
- Thiebot, J., Ito, K., Raclot, T., Poupart, T., Kato, A., Ropert-Coudert, Y., & Takahashi, A. (2016). On the significance of Antarctic jellyfish as food for Adélie penguins, as revealed by video-loggers. *Marine Biology*, *163*, 108. <https://doi.org/10.1007/s00227-016-2890-2>
- Thomas, D., & Dieckmann, G. (2010). *Sea ice* (2nd ed., 621 pp). Oxford, UK: Wiley-Blackwell.
- Vaughan, D. G., Comiso, J. C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., ... Zhang, T. (2013). Observations: cryosphere. *Climate Change*, *2103*, 317–382.
- Volkman, N. J., Presler, P., & Trivelpiece, W. Z. (1980). Diets of pygoscelid penguins at King George Island, Antarctica. *Condor*, *82*, 373–378. <https://doi.org/10.2307/1367558>
- Wang, S., Bailey, D., Lindsay, K., Moore, J. K., & Holland, M. (2014). Impact of sea ice on the marine iron cycle and phytoplankton productivity. *Biogeosciences*, *11*, 4713–4731. <https://doi.org/10.5194/bg-11-4713-2014>
- Watanabe, Y. Y., Ito, M., & Takahashi, A. (2014). Testing optimal foraging theory in a penguin – krill system. *Proceeding of the Royal Society*, *281*, 20132376. <https://doi.org/10.1098/rspb.2013.2376>
- Watanabe, Y. Y., & Takahashi, A. (2013). Linking animal-borne video to accelerometers reveals prey capture variability. *Proceedings of the National Academy of Sciences of the United States of America*, *110*, 2199–2204. <https://doi.org/10.1073/pnas.1216244110>
- Watanuki, Y., Kato, A., Naito, Y., Robertson, G., & Robinson, S. (1997). Diving and foraging behaviour of Adélie penguins in areas with and without fast sea-ice. *Polar Biology*, *17*, 296–304. <https://doi.org/10.1007/PL00013371>
- Watanuki, Y., Kato, A., Sato, K., Niizuma, Y., Bost, C.-A., Le Maho, Y., & Naito, Y. (2002). Parental mass change and food provisioning in Adélie penguins rearing chicks in colonies with contrasting sea-ice conditions. *Polar Biology*, *25*, 672–681.
- Widmann, M., Kato, A., Raymond, B., Angelier, F., Arthur, B., Chastel, O., ... Ropert-Coudert, Y. (2015). Habitat use and sex-specific foraging behaviour of Adélie penguins throughout the breeding season in Adélie Land. *East Antarctica. Movement Ecology*, *3*, 30. <https://doi.org/10.1186/s40462-015-0052-7>
- Wilson, R., & Wilson, M.-P. (1989). Tape: a package attachment technique for penguins. *Wildlife Society Bulletin*, *17*, 77–79.
- Woehler, E., & Johnstone, G. (1991). Status and conservation of the seabirds of the Australian Antarctic Territory. In J. P. Croxall (Ed.), *Seabird status and conservation: A supplement*. ICBP Technical Publication No. 11 (pp. 279–308). Cambridge, UK: International Council for Bird Preservation.
- Wood, S. (2006). *Generalized additive models: an introduction with R* (416 pp). Boca Raton, FL: Chapman and Hall/CRC.
- Xavier, J. C., Brandt, A., Ropert-Coudert, Y., Badhe, R., Gutt, J., Havermans, C., ... Sutherland, W. J. (2016). Future challenges in Southern Ocean ecology research. *Frontiers in Marine Science*, *3*, 94.
- Ydenberg, R., & Clark, C. (1989). Aerobiosis and anaerobiosis during diving by western grebes: an optimal foraging approach. *Journal of Theoretical Biology*, *139*, 437–449. [https://doi.org/10.1016/S0022-5193\(89\)80064-5](https://doi.org/10.1016/S0022-5193(89)80064-5)
- Youngflesh, C., Jenouvrier, S., Li, Y., Ji, R., Ainley, D. G., Ballard, G., ... Lynch, H. J. (2017). Circumpolar analysis of the Adélie Penguin reveals the importance of environmental variability in phenological mismatch. *Ecological Monographs*, *98*, 940–951.
- Zimmer, I., Wilson, R. P., Beaulieu, M., Ancel, A., Ropert-Coudert, Y., Kato, A., & Plötz, J. (2010). Dive efficiency in relation to depth in foraging emperor penguins. *Aquatic Biology*, *8*, 269–277. <https://doi.org/10.3354/ab00213>
- Zuur, A., Ieno, E., Walker, N., Saveliev, A., & Smith, G. (2009). *Mixed effects models and extensions in ecology with R* (574 pp.). New York, NY: Springer-Verlag. <https://doi.org/10.1007/978-0-387-87458-6>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Le Guen C, Kato A, Raymond B, et al. Reproductive performance and diving behaviour share a common sea-ice concentration optimum in Adélie penguins (*Pygoscelis adeliae*). *Glob Change Biol*. 2018;24:5304–5317. <https://doi.org/10.1111/gcb.14377>