



## High levels of fluoroalkyl substances and potential disruption of thyroid hormones in three gull species from South Western France



M. Sebastiano<sup>a,\*</sup>, W. Jouanneau<sup>a</sup>, P. Blévin<sup>a,b</sup>, F. Angelier<sup>a</sup>, C. Parenteau<sup>a</sup>, J. Gernigon<sup>c</sup>, J.C. Lemesle<sup>c</sup>, F. Robin<sup>c,d</sup>, P. Pardon<sup>e</sup>, H. Budzinski<sup>e</sup>, P. Labadie<sup>e</sup>, O. Chastel<sup>a</sup>

<sup>a</sup> Centre d'Etudes Biologiques de Chizé (CEBC), UMR 7372 CNRS-Univ. La Rochelle, France

<sup>b</sup> Akvaplan-niva AS, Fram Centre, NO-9296 Tromsø, Norway

<sup>c</sup> Réserve Naturelle de Lilleau des Niges, 17880, France


<sup>d</sup> Ligue pour la Protection des Oiseaux (LPO), 17300 Rochefort, France

<sup>e</sup> Univ. Bordeaux, CNRS, EPOC, EPHE, UMR 5805, F-33600 Pessac, France

### HIGHLIGHTS

- PFAS from three gull species from South western France were investigated.
- Positive relationships were found between exposure to PFAS and thyroid hormones.
- Great black-backed gulls document an increasing trend of plasma PFAS concentration.
- PFAS were negatively associated with the body condition of two of the studied species.
- Results on thyroid hormones and body condition were related to the sex of the birds.

### GRAPHICAL ABSTRACT



### ARTICLE INFO

#### Article history:

Received 30 September 2020

Received in revised form 8 December 2020

Accepted 16 December 2020

Available online 25 December 2020

Editor: Yolanda Picó

#### Keywords:

PFAS  
Seabirds  
South Western France  
Physiology

### ABSTRACT

Per- and poly-fluoroalkyl substances (PFAS) raised increasing concerns over the past years due to their persistence and global distribution. Understanding their occurrence in the environment and their disruptive effect on the physiology of humans and wildlife remains a major challenge in ecotoxicological studies. Here, we investigate the occurrence of several carboxylic and sulfonic PFAS in 105 individuals of three seabird species (27 great black-backed gull *Larus marinus*; 44 lesser black-backed gull *Larus fuscus graellsii*; and 34 European herring gull *Larus argentatus*) from South western France. We further estimated the relationship between plasma concentrations of PFAS and i) the body condition of the birds and ii) plasma concentrations of thyroid hormone triiodothyronine (TT3). We found that great and lesser black-backed gulls from South Western France are exposed to PFAS levels comparable to highly contaminated species from other geographical areas, although major emission sources (i.e. related to industrial activities) are absent in the region. We additionally found that PFAS are negatively associated with the body condition of the birds in two of the studied species, and that these results are sex-dependent. Finally, we found positive associations between exposure to PFAS and TT3 in the great black-backed gull, suggesting a potential disrupting mechanism of PFAS exposure. Although only three years of data have been collected, we investigated PFAS trend over the study period, and found that great black-backed gulls document an increasing trend of plasma PFAS concentration from 2016 to 2018. Because PFAS might have detrimental effects on birds, French seabird populations should be monitored since an increase of PFAS exposure may impact on population viability both in the short- and long-term.

© 2020 Elsevier B.V. All rights reserved.

\* Corresponding author.

E-mail address: [Manrico.Sebastiano@uantwerpen.be](mailto:Manrico.Sebastiano@uantwerpen.be) (M. Sebastiano).

## 1. Introduction

Per- and poly-fluoroalkyl substances (PFAS) raised increasing concerns over the past years due to their persistence and global distribution. Because of their high thermal and chemical stability, these synthetic substances have found an application in the manufacturing industry, mostly used as surfactants and additives (Buck et al., 2011), and have been widely produced over the past 50 years (Wang et al., 2017). Being extremely persistent in the environment, and due to their long-range transport via atmospheric and oceanic currents, they have been detected worldwide (Giesy and Kannan, 2001). Several studies have found PFAS to accumulate into living organisms (including invertebrates, fishes, amphibians, mammals, and birds) and to biomagnify through food webs (Kannan et al., 2005; Kelly et al., 2009; Simonnet-Laprade et al., 2019), and to date, PFAS exposure represents a global threat to human health and wildlife (Sunderland et al., 2019). Documenting their occurrence in the environment and understanding their disruptive effect on the physiology of humans and wildlife remains a major challenge.

Seabirds are long-lived apex predators generally exposed to high levels of environmental contaminants (Elliott and Elliott, 2013; Furness and Camphuysen, 1997), thus they prove particularly valuable to investigate PFAS accumulation in marine food webs especially in northern areas. High levels of PFOS have been found in plasma samples of several seabird species from the Arctic including ivory gulls *Pagophila eburnea* (average concentration of 31 ng/g, Lucia et al., 2017); glaucous gulls *Larus hyperboreus* (average concentration of 47 ng/g, Melnes et al., 2017; and of 134 ng/g, Verreault et al., 2005); in black-legged kittiwakes *Rissa tridactyla* from Svalbard (average concentration of 10.2 ng/g, Tartu et al., 2014); and in European shags *Phalacrocorax aristotelis* from Isle of May in Scotland (average concentration of 251 ng/g in females and 163 ng/g in males, Carravieri et al., 2020). High levels of PFOS were also found in egg samples of the European shag and common eider *Somateria mollissima* from Norway (average concentration of 36.8 ng/g and 37.4 ng/g, respectively, Herzke et al., 2009); in whole blood of the endangered lesser black backed gull *Larus fuscus* from Norway (average concentration of 33.5 ng/g, Bustnes et al., 2008a); and in several other seabird species. However, much work has been devoted to seabirds from the Arctic regions (i.e. considered a sink for environmental contaminants; Barrie et al., 1992; Braune et al., 2014; Wong et al., 2018), or in highly contaminated areas (e.g. China; Xie et al., 2013), while fewer studies have focused on areas with not-known sources of PFAS (i.e. Antarctica, Munoz et al., 2017b). In France, most studies examining PFAS occurrence and exposure in aquatic ecosystems focused on water, sediments, invertebrates, and fishes (Couderc et al., 2015; Fernandes et al., 2018; Munoz et al., 2019; Simonnet-Laprade et al., 2019). However, to the extent of our knowledge, no studies have been carried out on top predators including birds in this area, which may be exposed to concentrations of concern. It is therefore crucial to investigate PFAS exposure in French seabirds to document PFAS occurrence in marine biota and to provide early warning of its effects on their health status.

Over the past few years, there has been an increased body of evidence showing that PFAS may i) impact on adipogenesis thus with body condition (Tartu et al., 2014), and ii) disrupt several physiological traits of seabirds. For instance, previous work found that PFAS exposure is associated with lower levels of the stress hormone corticosterone (Tartu et al., 2014), higher oxidative stress (Costantini et al., 2019), longer telomeres (Blévin et al., 2017a; Sebastiano et al., 2020), and a higher metabolic rate (Blévin et al., 2017b). Further studies found PFAS to be associated with higher levels of the parental hormone prolactin and altered incubation behaviours (Blévin et al., 2020), lower hatching success (Tartu et al., 2014), and a higher survival rate (Sebastiano et al., 2020). Specifically, one way through which PFAS may impact on organism function is by disrupting hormonal mechanisms. Previous work provided evidence that PFAS have a strong affinity for proteins and are known to bind to the thyroid hormone transport protein transthyretin

(Ren et al., 2016; Weiss et al., 2009). In birds, the hypothalamic-pituitary-thyroid (HPT) axis controls the secretion of the thyroid hormone thyroxine (T4), which is then converted to triiodothyronine (T3), the active form of T4 (McNabb, 2007). Although Blévin et al. (2017b) found no association between PFAS exposure and thyroid hormones in adult black-legged kittiwakes, Braune et al. (2011) found a significant positive correlation between total triiodothyronine (TT3) levels and hepatic concentrations of PFAS in northern fulmars. Nøst et al. (2012) also found a positive association between PFAS levels and total thyroxine (TT4) in black-legged kittiwake and northern fulmar chicks, suggesting that PFAS may potentially act through an endocrine disrupting mechanism. More recently, Melnes et al. (2017) found that PFAS were positively associated with free triiodothyronine (FT3) in the glaucous gull. To date, further work is needed to understand the relationship between PFAS exposure and thyroid functioning in birds, especially considering that in birds, T3 and T4 are involved in a multitude of physiological pathways (McNabb, 2007). A disruption of thyroid hormone levels may be detrimental to development, behaviour, and reproduction (McNabb, 2007).

The Lilleau des Niges Natural Reserve is an important site for breeding, wintering, and migration of several bird species. It is located north of Ile de Ré, an island off the west coast of France, in front of La Rochelle, in the Bay of Biscay. By hosting several seabird species during the breeding season, this island offers a unique opportunity to investigate the occurrence of PFAS in a French seabird community. Although most previous studies have been carried out on a single species (Blévin et al., 2017b; Costantini et al., 2019; Melnes et al., 2017; Tartu et al., 2014), investigating several species simultaneously and from the same geographical area can help to better understand the mechanisms of exposure to PFAS and the potential physiological consequences of PFAS contamination. For instance, the Herring gull *Larus argentatus*, the lesser black-backed gull *Larus fuscus graellsii*, and the great black-backed gull *Larus marinus*, which breed sympatrically on the island, are characterized by different foraging and migratory strategies, thus potentially exposed to different concentrations of PFAS. The aims of this study were to i) assess to which extent French seabirds are contaminated by PFAS; ii) investigate the relationship between exposure to PFAS and body condition; and iii) determine the association between exposure to PFAS and plasma thyroid hormone T3 concentration in the three above mentioned seabirds from Ile de Ré. Data on PFAS occurrence and their potential adverse effects in seabirds from France are not yet available. To date, we are not aware of known point sources of PFAS in the region. But considering that diverse important rivers may discharge PFAS near the study area (Simonnet-Laprade et al., 2019), and that PFAS may reach and accumulate in remote areas due to their long-range oceanic and atmospheric transport (Munoz et al., 2019), we expect comparable PFAS concentrations with seabirds from the Arctic. In addition, if PFAS have a stimulating effect on thyroid hormone production as found in previous work (DeWitt, 2015; Liu et al., 2011; Nøst et al., 2012), we expect a positive association between PFAS and the concentration of thyroid hormones. Furthermore, although some PFAS are listed as POPs by the Stockholm Convention and their production has subsequently been reduced over the past years, studies investigating temporal trends of PFAS in tissues of wildlife are limited. Although our data have been solely collected over three years of study, we further aim to describe the temporal variation in blood concentration of PFAS from 2016 to 2018 in local seabirds.

## 2. Materials and methods

### 2.1. Sampling

Field work was performed in 2016, 2017, and 2018 at the Lilleau des Niges Natural Reserve (46° 13' 53" N, -1° 30' 22" W), managed by the Ligue pour la Protection des Oiseaux (LPO) located on the North side of Ile de Ré, France, as a part of a monitoring program for PFAS in the

region. A total of 108 breeding adult birds from three species were captured during the incubation stage on their nests using a nest trap. Because out of the 108 observations, three were coming from the same individuals sampled at different years, one or the other observation was randomly excluded to perform statistical analyses. Therefore, the final dataset included a total of 105 birds (European herring gull,  $n = 9$  in 2016,  $n = 16$  in 2017, and  $n = 9$  in 2018; lesser black-backed gull,  $n = 11$  in 2016,  $n = 17$  in 2017, and  $n = 16$  in 2018; great black-backed gull,  $n = 9$  in 2016,  $n = 7$  in 2017, and  $n = 11$  in 2018). After capture, 2 mL of blood was collected from the alar vein using a heparinized syringe and a 25 gauge needle. Blood was kept in a cold container and centrifuged for 10 min at  $8000 \times g$  at  $20^\circ\text{C}$  at the laboratory within a few hours after collection; plasma and red blood cells were kept frozen at  $-20^\circ\text{C}$  until laboratory analyses. Skull and tarsus were measured with an accuracy of 0.1 mm using a caliper. Wing length was also measured with an accuracy of 1 mm using a ruler, and birds were weighted to the nearest 5 g using a Pesola spring balance. Birds were sexed from red blood cells by polymerase chain reaction amplification (PCR) of part of two highly conserved genes (CHD) of sexual chromosomes. Briefly, DNA was extracted from erythrocytes and the sex was determined by molecular sexing based on PCR amplification of the CHD gene as described in Fridolfsson and Ellegren (1999). Amplification was performed in 20  $\mu\text{L}$  final volume with a Eppendorf Mastercycler using 0.5 U Taq DNA polymerase, 200  $\mu\text{M}$  dNTPs, 10 mM Tris-HCl pH 8.3, 50 mM KCl, 1.5 mM  $\text{MgCl}_2$  and 0.4  $\mu\text{M}$  of primers 2550F (5'-GTTACTGATTCGTCTACGAGA-3') and 2718R (5'-ATTGAAATGATCCAGTGCTTG-3'). Female birds may deposit a significant amount of PFAS into their eggs. Therefore, to minimize the variation due to PFAS deposition in eggs, we have only sampled individuals having either two (17/105, 16% of birds) or three eggs (87/105, 83% of birds) except one sampled females that only laid one egg (1/105, 1%). Preliminary statistical analyses were carried out to test whether females with two (8/52, 15% of females) or three eggs (43/52, 83%) had different concentrations of PFAS. However, linear models showed that for any PFAS, concentrations were similar between females that laid two or three eggs (all  $t < 1.14$ , all  $P > 0.26$ ), thus clutch size was not further included in the statistical analyses.

## 2.2. PFAS analyses

A total of 14 PFAS were analysed in each plasma sample, including eight carboxylates: branched- (Br-PFOA) and linear-perfluorooctanoate (L-PFOA), perfluorononanoate (PFNA), perfluorodecanoate (PFDA), perfluoroundecanoate (PFUnDA), perfluorododecanoate (PFDoDA), perfluorotridecanoate (PFTrDA), perfluorotetradecanoate (PFTeDA); and six sulfonates: perfluorohexanesulfonate (PFHxS), branched- (Br-PFHps) and linear-perfluoroheptanesulfonate (L-PFHps), branched perfluorooctanesulfonate (Br-PFOS), linear perfluorooctanesulfonate (L-PFOS), and perfluorooctanesulfonamide (FOSA). Analytical standards of native PFAS along with a series of  $10^{13}\text{C}$ ,  $18\text{O}$  or D mass-labelled internal standards used for quantification purposes were supplied by Wellington laboratories. All reagents were analytical grade or equivalent (see Munoz et al. (2017b) for full details). Briefly, in a 2 mL polypropylene Eppendorf tubes, a 25  $\mu\text{L}$  aliquot of plasma was weighed (~25 mg) and internal standards (ISs) were subsequently added under gravimetric control (~15 mg of a 1  $\mu\text{g}/\mu\text{L}$  IS mixture prepared in methanol). Following protein precipitation with 100  $\mu\text{L}$  of acetonitrile (ACN), extracts were centrifuged for 10 min at  $24,000 \times g$  at  $20^\circ\text{C}$ . The supernatant was then transferred to 2 mL polypropylene centrifuge tubes (0.22  $\mu\text{m}$  nylon filter). After centrifugation for 3 min at  $7000 \times g$  at  $20^\circ\text{C}$ , extracts were transferred to 2 mL auto sampler glass vials and diluted with 675  $\mu\text{L}$  of HPLC-water. Extracts were briefly vortexed and then processed using an Agilent Technologies (Massy, France) on-line SPE platform which comprises a standard auto sampler (1260 Infinity ALS), a quaternary pump (1260 Infinity Quaternary Pump VL), a switch valve (1200 2 Position/6 Port Valve) and an on-line SPE column support (1200 6 Position Selection Valve), which

were all automatically controlled via the Acquisition module of the Agilent Mass Hunter software as previously done (Munoz et al., 2017b). HPLC-water aliquots were run between each seabird plasma sample to eliminate any cross-contamination. Note that on-line extraction was performed with Waters Oasis HLB on-Line SPE columns ( $2 \times 10$  mm,  $\text{dp} = 25\text{--}35 \mu\text{m}$ ) while analyte separation was carried out using an Agilent  $\text{C}_{18}$  Poroshell analytical column ( $2.1 \times 100$  mm, 2.7  $\mu\text{m}$ ).

## 2.3. Quality assurance/quality control (QA/QC)

When analytes were detected in blanks, blank correction was performed and a limit of reporting (LOR) was defined as three times the maximum blank signal divided by the average mass of plasma used for analysis. A limit of detection (LOD) was also defined as the concentration yielding a signal to noise ratio of 3 in spiked plasma samples. Because laboratory analyses were performed in different years, a unique left-censoring threshold was set for each analyte, i.e. the maximum between LORs and LODs, all years combined. For those PFAS with concentrations below this threshold in less than 30% of samples, left-censored data were arbitrarily replaced with  $\frac{1}{2} \times \text{LOR}$  or  $\text{LOD}$  to enable statistical analyses. Therefore, ten PFAS (PFNA, PFDA, PFUnDA, PFDoDA, PFTrDA, PFTeDA, PFHxS, L-PFHps (hereafter PFHpS), Br-PFOS, and L-PFOS) could be further investigated (i.e. other analytes were excluded from statistical analyses). LORs, LODs and detection frequencies are presented in the supplementary information (Table S1). For each sample batch (20 samples), several QA/QC points were assessed by analyzing: i) two procedural blanks consisting of 25  $\mu\text{L}$  of HPLC-water that went through the entire analytical procedure; ii) one human serum standard reference material (NIST SRM 1957, trueness assessment); iii) replicate spiked chicken plasma samples (target analytes added jointly with mass-labelled ISs at the beginning of the preparation procedure at 2 ng/g each, accuracy assessment); and iv) HPLC-water samples spiked at 2 ng/g, accuracy assessment) as previously described (Munoz et al., 2017b). Procedural blanks showed very limited contamination. The analysis of NIST SRM 1957 gave satisfactory results, i.e. within the specified uncertainty interval. For those compounds with a reference concentration (i.e. PFHpA, PFOA, PFNA, PFDA, PFUnDA, PFHxS and PFOS), levels deviated between 2 and 20% from the reference concentration (except for FOSA, which deviated 36%).

## 2.4. Thyroid hormone analyses

TT3 was determined by radioimmunoassay. Briefly, 25  $\mu\text{L}$  of plasma was incubated for 24 h at  $4^\circ\text{C}$  with a known concentration (10,000 cpm) of T3 marked with the radioisotope Iodine-125 ( $\text{T3-}^{125}\text{I}$ , Perkin Elmer, US, reference: NEX110X100UC) and an antibody *Ab* (polyclonal rabbit antiserum, Sigma-Aldrich, US, reference: T-2777). Because *Ab* is available in a limited concentration, T3 and  $\text{T3-}^{125}\text{I}$  compete for *Ab*, to which they bind. Therefore, after incubation, there is a bound fraction (T3 and  $\text{T3-}^{125}\text{I}$  bound to *Ab*) and a free fraction (T3 and  $\text{T3-}^{125}\text{I}$  unbound to *Ab*), which are separated by adding a sheep anti-rabbit antibody (whole anti-serum anti rabbit IgG produced in sheep), incubated for 12 h at  $4^\circ\text{C}$  followed by centrifugation at  $4300 \times g$  at  $18\text{--}20^\circ\text{C}$  for 45 min. The bound fraction is then counted with a wizard 2 gamma counter (Perkin Elmer, US). Pooled plasma of diverse gull samples were serially diluted and produced a dose-response curve parallel to the T3 standard curve. The lowest TT3 detectable concentration was 0.07 ng/ml (LOD). Samples below this limit ( $n = 3$ ) were replaced with a value equal to  $\frac{1}{2} \times \text{LOD}$  to enable statistical analyses. All samples were run in duplicates. Samples that had a coefficient of variation above 15% and could not be done in triplicates due to low plasma volume were not included in statistical analyses ( $n = 11$ ). An additional measurement of TT3 was excluded from statistical analyses since it was considered an outlier (the measurement exceeded the mean  $\pm 3$  times the standard deviation and was highly influential in statistical analyses). Therefore, for a total of 58 samples ( $n = 6$  in 2016 and  $n = 12$  in 2017 for the European herring gull;  $n = 10$  in 2016 and  $n = 14$  in

2017 in lesser black-backed gull;  $n = 8$  in 2016 and  $n = 8$  in 2017 in great black-backed gull), both TT3 and PFAS data were available. The intra-assay coefficient of variation was 9.73%, while the inter-assay coefficient of variation amounted to 15.13%.

## 2.5. Statistical analyses

After PFAS data were log-transformed to reduce the influence of extreme values (see below), linear models were used to investigate differences in PFAS concentrations among species and between genders (in samples collected from 2016 to 2018). In each model, each PFAS was considered as a dependent variable while the factors *Species*, *Sex*, and their interaction were considered as predictors. To simultaneously investigate species specific temporal trends in PFAS exposure from 2016 to 2018, the *Year*, the factor *Species*, and their interaction were also included in the model as explanatory variables. We used a similar model to test the difference in body condition between sexes. Briefly, the body condition has been calculated using the body mass adjusted by a linear body measurement (i.e. skull length) using the formula described in Peig and Green (2009).

Linear models were additionally used to study the association between TT3 and PFAS (in samples collected in 2016 and 2017). In these models, a three-way interaction between PFAS, the factors *Species* and *Sex*, was used to investigate sex-related responses to PFAS exposure. These models additionally included the *Year* (as a factor, to control for

the temporal variation in TT3 and PFAS), and *body condition* (as a covariate, to control for the individual condition of birds). A similar model was built to test the association between the body condition and PFAS in all samples collected from 2016 to 2018.

All PFAS concentrations (except PFTeDA, which was normally distributed and assumptions listed below were respected without data transformation), were log-transformed when testing for time trends and when testing for inter-species and between sex differences. All PFAS concentrations were log-transformed when testing for the association between either TT3 or body condition and PFAS. Data transformation was done to meet model assumptions as homoscedasticity and normality of residuals, further confirmed by visually inspecting Q-Q plots. All data transformation and violation of models' assumptions are reported throughout the manuscript. Statistical significance was set to  $\alpha = 0.05$  and 95% confidence intervals were used during data processing and data visualization. All statistical analyses were performed using R version 3.5.2.


## 3. Results

PFAS used in statistical analyses were detected in all samples (Table S1) and their concentrations are summarized in Table 1 and Fig. 1. L-PFOS was the most abundant, followed by the odd-chain carboxylates PFTTrDA and PFUnDA. Linear models showed statistically significant differences among species for all carboxylic and sulfonic PFAS (all  $F > 5.15$ , all  $P < 0.01$ ), and all statistical outputs and post-hoc

**Table 1**

PFAS concentration (ng/g of ww) in females and males of the three seabird species from Ile de Re. *P*-values refer to the difference in PFAS between females and males. Significant *P*-values are bolded. SE refers to standard errors.

Herring gull <i>Larus argentatus</i>					
	Females (n = 18)		Males (n = 16)		<i>P</i> -value
	Mean ± SE	Median (range)	Mean ± SE	Median (range)	
PFNA	1.16 ± 0.11	1.06 (0.36, 2.16)	1.68 ± 0.17	1.63 (0.67, 2.85)	0.07
PFDA	1.16 ± 0.19	0.94 (0.38, 4.01)	1.98 ± 0.30	1.79 (0.69, 6.04)	<b>0.01</b>
PFUnDA	1.23 ± 0.17	1.01 (0.47, 3.45)	2.43 ± 0.34	2.17 (0.61, 6.49)	<b>&lt;0.001</b>
PFDoDA	0.95 ± 0.16	0.9 (0.3, 3.22)	1.82 ± 0.32	1.56 (0.55, 6.17)	<b>&lt;0.001</b>
PFTTrDA	1.76 ± 0.27	1.42 (0.55, 5.25)	3.03 ± 0.29	3.05 (1.04, 4.81)	<b>&lt;0.001</b>
PFTeDA	0.61 ± 0.07	0.61 (0.25, 1.23)	1.23 ± 0.16	1.03 (0.33, 2.8)	<b>0.04</b>
PFHxS	2.26 ± 0.21	2.16 (0.88, 3.5)	2.29 ± 0.36	1.89 (0.79, 6.92)	0.99
PFHpS	0.27 ± 0.02	0.25 (0.11, 0.5)	0.49 ± 0.09	0.38 (0.11, 1.63)	<b>0.045</b>
Br-PFOS	2.03 ± 0.34	1.71 (0.86, 6.8)	3.95 ± 0.45	3.75 (1.07, 7.9)	<b>&lt;0.001</b>
L-PFOS	13.82 ± 2.0	11.64 (5.57, 37.04)	30.44 ± 4.31	28.44 (6.6, 62.52)	<b>&lt;0.001</b>
Lesser black-backed gull <i>Larus fuscus graellsii</i>					
	Females (n = 20)		Males (n = 24)		<i>P</i> -value
	Mean ± SE	Median (range)	Mean ± SE	Median (range)	
PFNA	1.14 ± 0.09	1.11 (0.46, 1.83)	2.04 ± 0.1	2.06 (0.95, 3.36)	<b>&lt;0.001</b>
PFDA	1.1 ± 0.08	1.15 (0.54, 1.64)	2.63 ± 0.15	2.58 (1.17, 4.16)	<b>&lt;0.001</b>
PFUnDA	1.88 ± 0.16	1.69 (0.66, 3.28)	5 ± 0.29	4.89 (1.98, 8.7)	<b>&lt;0.001</b>
PFDoDA	1.02 ± 0.07	1.02 (0.39, 1.87)	3.06 ± 0.2	2.86 (1.75, 5.52)	<b>&lt;0.001</b>
PFTTrDA	1.97 ± 0.12	2.06 (0.66, 2.91)	6.38 ± 0.44	5.74 (3.53, 11.66)	<b>&lt;0.001</b>
PFTeDA	0.66 ± 0.07	0.61 (0.13, 1.39)	2.23 ± 0.2	1.87 (1.1, 4.35)	<b>&lt;0.001</b>
PFHxS	2.18 ± 0.25	1.95 (1.12, 6.01)	2.56 ± 0.26	2.24 (1.08, 5.56)	0.94
PFHpS	0.37 ± 0.03	0.35 (0.16, 0.82)	0.79 ± 0.07	0.69 (0.34, 1.84)	<b>&lt;0.001</b>
Br-PFOS	1.86 ± 0.19	1.96 (0.58, 3.15)	5.06 ± 0.5	4.41 (2.38, 13.67)	<b>&lt;0.001</b>
L-PFOS	21.09 ± 2.06	22.76 (5.74, 34.49)	60.23 ± 4.44	54.68 (26.51, 119.69)	<b>&lt;0.001</b>
Great black-backed gull <i>Larus marinus</i>					
	Females (n = 14)		Males (n = 13)		<i>P</i> -value
	Mean ± SE	Median (range)	Mean ± SE	Median (range)	
PFNA	0.59 ± 0.06	0.59 (0.24, 0.95)	1.94 ± 0.29	1.72 (0.89, 4.85)	<b>&lt;0.001</b>
PFDA	1.41 ± 0.23	1.13 (0.32, 3.55)	3.83 ± 0.58	3.34 (1.52, 7.2)	<b>&lt;0.001</b>
PFUnDA	2.42 ± 0.35	2.23 (0.49, 4.93)	4.67 ± 0.65	4.08 (2.62, 11.46)	<b>0.01</b>
PFDoDA	1.69 ± 0.24	1.63 (0.32, 3.12)	2.79 ± 0.29	2.37 (1.68, 5)	<b>0.04</b>
PFTTrDA	4.04 ± 0.72	3.55 (0.68, 9.61)	6.87 ± 0.83	5.79 (3.85, 14.39)	<b>0.02</b>
PFTeDA	0.96 ± 0.17	0.92 (0.13, 2.4)	1.94 ± 0.32	1.42 (0.91, 4.17)	<b>0.03</b>
PFHxS	1.2 ± 0.11	1.16 (0.62, 2.06)	2.01 ± 0.32	1.58 (0.93, 5.33)	0.16
PFHpS	0.2 ± 0.03	0.15 (0.09, 0.47)	0.63 ± 0.21	0.42 (0.22, 3.01)	<b>&lt;0.001</b>
Br-PFOS	1.32 ± 0.17	1.1 (0.39, 2.53)	4.21 ± 0.89	3.21 (1.85, 13.01)	<b>&lt;0.001</b>
L-PFOS	13.45 ± 1.52	10.53 (5.18, 25.92)	46.55 ± 13.74	27.62 (16.71, 194.72)	<b>&lt;0.001</b>



**Fig. 1.** Plasma concentrations of PFAS (expressed as ng/g of ww) in the three seabird species from Ile de Re. Statistically significant differences are indicated by the asterisk; \*, \*\*, \*\*\*, indicate a  $P$ -value  $< 0.05$ ,  $< 0.01$ , and  $< 0.001$ , respectively.

differences can be found in Table S2. Among carboxylates, PFNA levels were higher in herring gulls and lesser black-backed gulls than great black-backed gulls (both  $P < 0.05$ , Fig. 1), PFDA levels were higher in great black-backed gulls than herring gulls ( $P < 0.01$ , Fig. 1), PFUnA, PFDODA, PFTTrDA, and PFTeDA levels were higher in great and lesser black-backed gulls than herring gulls (all  $P < 0.01$ , Fig. 1), and PFTTrDA levels were also higher in great than lesser black-backed gulls ( $P < 0.05$ , Fig. 1). Among sulfonic acids, PFHxS was higher in both lesser black-backed and herring gulls than great black-backed gulls (both  $P < 0.01$ , Fig. 1), PFHpS and L-PFOS were highest in lesser black-backed gulls (all  $P < 0.001$ , Fig. 1), and Br-PFOS was higher in lesser than great black-backed gulls ( $P < 0.05$ , Fig. 1).

All carboxylates showed significantly higher concentrations in males than in females for all species (all  $P < 0.05$ , except for PFNA in the herring gull, for which  $P = 0.07$ ; Table 1). Among sulfonates, PFHxS showed similar concentrations between females and males in all species (all  $P > 0.16$ , Table 1), while PFHpS, Br- and L-PFOS showed significantly higher concentrations in males than females (all  $P < 0.05$ , Table 1). In all three species, there was no difference in the body condition between males and females (all  $t < 1.27$ , all  $P > 0.80$ ). Finally, TT3 levels were similar between sexes in all three species ( $t = 2.76$ ,  $P = 0.08$ ).

In great black backed gulls, TT3 was positively associated with PFUnDA, PFDODA, PFTTrDA, PFTeDA and Br-PFOS in females (all  $t > 2.10$ , all  $P \leq 0.04$ ; Fig. 2a–d, f, Table S3), while TT3 was negatively associated with PFHxS in males ( $t = -2.69$ ,  $P = 0.01$ ; Fig. 2e, Table S3). There was no association between TT3 and any PFAS in herring gulls and lesser black-backed gulls (all  $t \leq 0.79$ , all  $P \geq 0.43$ , Table S3).


In great black backed gull females, increasing levels of PFNA, PFDA, PFHxS, PFHpS, Br- and L-PFOS were associated with a reduced body condition (all  $t \leq -2.19$ , all  $P \leq 0.03$ ; Figs. S1, S2, Table S4), while increasing levels of PFNA and PFDA were associated with a reduced body condition (both  $t \leq -2.18$ , both  $P = 0.03$ ; Fig. S1, Table S4) in lesser black-backed gull males, but not females.

In great black backed gulls, there was a significant or marginally-significant increase in all PFAS from 2016 to 2018 (all  $t \geq 1.93$  and all  $P \leq 0.056$ ; Figs. 3 and 4, Table S5), while lesser black backed gulls showed an increase in PFTeDA and PFHxS, and a marginally significant decrease in PFUnDA levels from 2016 to 2018 (all  $t \geq 1.98$ , all  $P \leq 0.05$ ; Figs. 3, 4, Table S5). All PFAS showed similar concentrations among years in the herring gulls except for PFTTrDA, which increased from 2016 to 2018 ( $t = 2.30$ ,  $P = 0.02$ ; Fig. 3, Table S5). Body condition did not change in any of the species from one year to another (all  $t < 1.14$ , all  $P > 0.26$ ). TT3 levels remained similar between 2016 and 2017 in all three species (all  $t < 1.88$ , all  $P > 0.42$ ).

#### 4. Discussion

Our study is the first to provide evidence that although not-known point sources of emission are present in the region, several PFAS were detected in seabird species from South Western France. Great and lesser black-backed gulls show that both plasma carboxylate and sulfonate concentrations are comparable to highly contaminated seabird species from Arctic regions, while herring gulls are exposed to relatively lower levels of PFAS. We found that PFAS are negatively associated with the body condition of the birds. Furthermore, TT3 levels were associated with several PFAS in a contrasted manner between sexes in the great black-backed gull, suggesting a potential disrupting mechanism of PFAS exposure. Finally, the great black-backed gulls documented an increasing trend of plasma PFAS concentration from 2016 to 2018.

Our results show that among carboxylates, PFUnDA and PFTTrDA are the most abundant congener in all three species, a pattern that is commonly found in seabird species (Bustnes et al., 2008b; Melnes et al., 2017; Tartu et al., 2014). Because of the strong winds and oceanic currents that characterize the Atlantic Ocean, and considering that, to the best of our knowledge, there are no point sources of PFAS in the region, perfluorinated compounds should occur at a lower concentration than in the Mediterranean and the Arctic regions, considered as sinks for pollutants (Danovaro, 2003; Wong et al., 2018). However, the levels of carboxylates found in this study (ranging from a median of 0.6 ng/g of PFTeDA in Herring gulls to 5.8 ng/g of PFTTrDA in great black-backed gulls) are similar to those reported for glaucous gulls (ranging from a median of 0.1 ng/g of PFOA to 3.8 ng/g of PFUnDA, Melnes et al., 2017) and lesser black-backed gulls (ranging from a median of 0.2 ng/g of PFDODA to 5.9 ng/g of PFDA, Bustnes et al., 2008a) from Arctic regions, but lower than those found in other species (e.g. in black-legged kittiwakes from Svalbard; ranging from a mean of 1.0 ng/g of PFNA to 18.2 ng/g of PFTTrDA, Tartu et al., 2014). One possible explanation may be related to the continental input of PFAS through the Gironde, Loire, and Charente estuaries (Munoz et al., 2019; Munoz et al., 2017a; Munsch et al., 2019), which may have contributed to the observed concentrations. Except for PFNA (which levels were higher in herring gulls and lesser black-backed gulls than great black-backed gulls), most carboxylates were higher in lesser and great black backed gulls than herring gulls. Although we cannot exclude that these three species differ in their ability to excrete PFAS from their body, our results suggest that the differences in exposure likely depend on the trophic niche occupied by the species. For instance, a recent study on PFAS in six seabird species from the Arctic regions showed that predatory birds (e.g. great skua *Stercorarius skua*, 44.8 ng/g of sumPFAS) showed the highest contaminant load compared with species from a lower trophic level (e.g. common eider *Somateria mollissima*, 1.3 ng/g of sumPFAS; Haarr et al.,




**Fig. 2.** Relationship between the concentration of the thyroid hormone TT3 (expressed as ng/mL) and log-transformed carboxylic (C-11 to C-14, panel a to d, respectively), and sulfonic PFAS (PFHxS and Br-PFOS, panel e and f, respectively) of the three seabird species from Ile de Re. The solid line represents the trend while curved dashed lines represent 95% confidence intervals. Shapes of data points and line colours are used to distinguish the three species as explained in the figure legend. Only significant trends are shown. Data refer to the period 2016–2017 for which both TT3 and PFAS were available ( $n = 58$ ).

2018). Great black-backed gulls feed on higher trophic level preys and mainly forage along the shore (Maynard and Davoren, 2020), while lesser black-backed and herring gulls are known to have a generalist diet which also includes food items from both terrestrial and marine origin (Corman et al., 2016; Maynard and Davoren, 2020).

Among sulfonates, L-PFOS was the most abundant, followed by Br-PFOS, PFHxS, and PFHpS and all occurred at very high concentrations. For instance, PFHxS ranged from a median value of 1.2 ng/g in great black-backed gulls to 2.2 ng/g in lesser black-backed gulls, while other studies from highly contaminated areas reported lower PFHxS plasma concentrations (a median below 1 ng/g in lesser black-backed gulls, Bustnes et al., 2008b; a median below 0.7 in glaucous gulls, Melnes et al., 2017; all samples below 0.2 ng/g in black-legged kittiwakes,

Tartu et al., 2014). Furthermore, L-PFOS ranged from a median of 11.6 ng/g in herring gull females to a median of 54.7 ng/g in lesser black-backed gull males. Thus, lesser black-backed gulls in our study showed very high sulfonate levels, even higher than Norwegian populations (a median of 40 ng/g of PFOS in males, Bustnes et al., 2008b), and in this species, sulfonate levels are significantly higher than those observed in great black-backed gulls. In addition, plasma concentrations of most PFAS showed significantly higher levels in males than in females, but this difference was not related to the body condition of the birds, with males showing a similar body condition than females. Because females transfer contaminants in the eggs, it is thus possible that females have lower levels of circulating PFAS in plasma. However, previous work pointed out contrasting results between PFAS in eggs




**Fig. 3.** Error bar plots mean  $\pm$  standard error of carboxylate concentrations in 2016, 2017, and 2018 in the seabird species from Ile de Re. The shapes and the three different grades of grey are used to distinguish the three species as explained in the figure legend. Only significant trends are shown. Dashed lines indicate a trend close to significance.

and plasma, suggesting that the extent of PFAS transfer to the eggs may significantly vary among the studied species (Bustnes et al., 2008a; Herzke et al., 2009; Verreault et al., 2005). Given that in this study we did not analyse PFAS levels in eggs, it is not possible to clarify whether the species differ in terms of PFAS excreted in eggs. Although Verreault et al. (2006) found that the contaminant content in gull eggs fluctuated irrespectively of the laying order, other work suggests that the majority of PFAS are found in the first or the first two eggs, while negligible concentrations of PFAS are found in the third egg, as previously shown in Audouin's gulls *Larus audouinii* (Vicente et al., 2015). Our results showed that females that laid three eggs had similar PFAS concentrations than females that laid two eggs, therefore our results should not be affected by the difference in PFAS deposition in eggs.

Because of the higher bioaccumulative properties and biomagnification of longer-chain PFAS (Boisvert et al., 2019; Simonnet-Laprade

et al., 2019), these compounds tend to occur at higher concentrations in wildlife tissues (Conder et al., 2008; Muir et al., 2019; Muir and de Wit, 2010). Therefore, individuals feeding at a higher trophic position are likely to be exposed to higher concentrations of long-chained carboxylates. Previous work also showed that longer chained PFAS are more likely to induce adverse health effects in seabirds compared to shorter chained PFAS. For instance, negative associations between PFAS and baseline corticosterone in black-legged kittiwakes were only found for PFTrDA and PFTeDA (Tartu et al., 2014), while such association was not found for shorter-chain PFAS. Additional work on the same bird population found higher protein oxidative damage in those birds having higher concentrations of PFDoDA, PFTrDA and PFTeDA (Costantini et al., 2019). Similarly, previous work found a positive association between PFTrDA and metabolic rate in the same species (Blévin et al., 2017b). These results were further corroborated by experimental work on rat *Rattus* sp. cell cultures, showing that the cytotoxicity of PFAS increases



**Fig. 4.** Error bar plots mean  $\pm$  standard error of sulfonate concentrations in 2016, 2017, and 2018 in the seabird species from Ile de Re. The shapes and the three different grades of grey are used to distinguish the three species as explained in the figure legend. Only significant trends are shown. Dashed lines indicate a trend close to significance.

with increasing carbon chain length (Berntsen et al., 2017), and that comparing molecules with a similar chain length, a sulfonate functional group may lead to greater toxicity than a carboxyl group (Berntsen et al., 2017).

Our results are thus of particular interest as they suggest that birds feeding at a higher trophic position and showing a more marine diet (i.e. great black-backed gulls) should also be exposed to greater toxicological risks from carboxylates, while birds with a more generalist diet (i.e. lesser black-backed gulls) can be exposed to higher levels of sulfonates. This is likely the reason why we found associations between TT3 and PFAS only in great black-backed gulls, while no significant associations have been found in herring gulls and lesser black-backed gulls. Interestingly, our results were dependent on the sex of the birds. Indeed, although we found that PFHxS was associated with TT3 in great black-backed gull males, all other associations were found in females only. Despite the lower concentration of circulating plasma PFAS levels, females can deposit PFAS in eggs therefore we cannot be certain that females were exposed to lower PFAS concentrations than males, and further work including other tissues (e.g. liver or muscle) would clarify whether PFAS intake differs between the sexes of sampled birds. Despite the absolute concentrations to which they are exposed, a possible explanation for the results in females may rely on the fact that incubation can be extremely costly for female birds (Hanssen et al., 2005), thus they may be more susceptible to PFAS exposure. Previous work in birds suggest a modulation of thyroid function induced by exposure to various environmental contaminants. Smits et al. (2002) reported decreased TT3 levels in American kestrels *Falco sparverius* experimentally exposed to PCBs, while Verreault et al. (2004) reported a decrease in T4:T3 ratio in the glaucous gull. Similarly, exposure to organochlorines was associated with reduced TT3 in kittiwakes (Blévin et al., 2017b), and with reduced T3 and T4 in glaucous gulls (Melnes et al., 2017; Verreault et al., 2004). However, specifically related to PFAS, further work on seabirds found a positive association with thyroid functioning (i.e. between PFOS and

TT3 in glaucous gulls, Melnes et al., 2017; between several PFAS and TT4 in black-legged kittiwakes and northern fulmars, Nøst et al., 2012). Although being conducted in fish, an experimental approach showed that exposure to PFOS in zebra fish (*Danio rerio*) led to increased thyroid hormones secretion (Liu et al., 2011). Thus, our results on great black backed gulls are in line with previous studies. In this study, increasing TT3 levels in this species were found with increasing concentrations of longer chain PFAS (PFUnDA, PFDoDA, PFTrDA, and PFTeDA) and Br-PFOS. This suggests that despite carboxylates and sulfonates are functionally different, their effect on TT3 is similar. However, this was not the case for PFHxS, which showed a decrease in TT3 levels with increasing concentrations. PFHxS is highly toxic and causes thyroid disruption by lowering thyroid hormone levels in rats (Ramhøj et al., 2020), but it remains unclear why this effect was only found in great black-backed gull males. Indeed, females showed similar levels than males, and the other species exhibited higher PFHxS concentrations than those found in great black backed gulls, thus this result would strongly benefit from experimental support. Because the avian thyroid gland secretes almost exclusively T4 (Darras et al., 2006), most T3 is derived from the deiodination of T4 (Darras et al., 2006). A possible explanation is that in great black-backed gulls, exposure to PFHxS may negatively impact either the transport of T4 (by reducing the activity of serum binding proteins) or deiodination processes. It is therefore strongly warranted to supplement in vitro experiments to verify the effect of PFHxS on T4 transformation. Our results do not provide evidence for a causal relationship PFAS exposure and circulating thyroid hormones. But the contrasting results found between sexes strongly call for further work to experimentally investigate the effect of PFAS exposure on thyroid functioning of birds.

Sex-related differences were also found while investigating the relationship between exposure to PFAS and the birds' body condition. In female great black-backed gulls, we found that increasing concentrations of PFNA, PFDA, and all four sulfonates were

negatively associated with body condition, while in lesser black-backed gulls, a similar negative relationship between PFNA, PFDA, and body condition was only found in males. Previous work in humans and various animal models found that exposure to certain PFAS is suspected to disrupt fatty acids metabolism and promote adipogenesis (Cheng et al., 2016; Wan et al., 2012; Xu et al., 2016; Yeung et al., 2007). More specifically, these changes in lipid content are related to the capacity of PFAS to alter the expression of genes involved in the metabolism of lipids and fatty acids (Jacobsen et al., 2018; Wan et al., 2012). To date, work on the effect of PFAS on lipid metabolism and body condition in birds remains extremely limited. The negative association we found is in contrast with a previous study on black-legged kittiwakes, showing that PFNA was positively associated with body condition in males (Tartu et al., 2014). One possible explanation for the negative relationship found in females may be related to the ability to deposit PFAS into the eggs. For instance, great black-backed gull females in a better body condition may be more efficient in eliminating PFAS through egg-deposition, although evidences to support this statement are lacking. However, this would not explain why in a closely related species (i.e. lesser black-backed gull) a similar association has been found in males. Our study results are novel but emphasize the need to experimentally investigate the potential association between exposure to PFAS and body condition in birds.

Finally, not only does our work provide evidence of high PFAS levels in seabirds from metropolitan France, but our results clearly suggest increasing blood concentration of most PFAS over a relatively short period of time (i.e. from 2016 to 2018) in great and lesser black-backed gulls. A previous study showed that birds caught later on over the breeding season had lower concentration of PFAS (Bustnes et al., 2008a). However, all birds included in this study were sampled during the same period of the year (difference of a few days from one year to another), and, more importantly, all birds were sampled while incubating eggs, thus during the same reproductive state, which should not affect the results. Additionally, we cannot exclude that we unintentionally captured older birds in more recent years (assuming that PFAS levels increase with age in these species). However, considering that all individuals included in this study were adults, and assuming that PFAS concentrations in birds reach a steady level relatively early in life as previously shown for organochlorines (Bustnes et al., 2003), these trends should not be affected by the age of the bird. Information on temporal trends of PFAS in birds' tissues in recent years are scarce and do not exhibit any overall trend (Jouanneau et al., 2020; Land et al., 2018; Muir et al., 2019; Sun et al., 2019). Thus, although only three years of data could be included in the present study, our results provide valuable information on PFAS trends in South western France. Further work including several years of study is strongly warranted to corroborate these findings.

## 5. Conclusions

Our study provides the first evidence of the presence of high levels of PFAS in seabirds from South western France. Despite some PFAS showed similar levels of other seabird species, L-PFOS and some other PFAS showed either comparable or higher levels than highly contaminated seabird species, and may therefore pose a threat to long-lived seabirds. This hypothesis is further corroborated by our results showing an association between PFAS and the level of the thyroid hormone TT3. Similarly, we provide evidence that PFAS may interfere with lipid accumulation and body condition in birds, and we call for further work to experimentally test this hypothesis. Work on species-specific mechanisms of contaminant excretion and susceptibility to PFAS exposure would prove useful to understand the consequences of PFAS exposure in different species. Our results also document an increase in blood PFAS concentrations

over time, particularly in great black-backed gulls, suggesting that PFAS concentrations may also be increasing in the investigated species. Because PFAS have detrimental effects on birds, these and other seabird populations should be monitored as an increase of PFAS exposure may impact on population viability both in the short- and long-term.

## CRedit authorship contribution statement

**M. Sebastiano:** Writing – original draft, Resources, Investigation. **W. Jouanneau:** Writing – review & editing, Resources, Investigation. **P. Blévin:** Writing – review & editing, Resources, Investigation. **F. Angelier:** Writing – review & editing, Resources, Investigation. **C. Parenteau:** Resources, Formal analysis. **J. Gernigon:** Resources, Investigation. **J.C. Lemesle:** Resources, Investigation. **F. Robin:** Resources, Investigation. **P. Pardon:** Resources, Formal analysis. **H. Budzinski:** Writing – review & editing, Resources, Formal analysis. **P. Labadie:** Writing – review & editing, Resources, Formal analysis. **O. Chastel:** Writing – review & editing, Resources, Investigation, Supervision.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

This work was funded by the Région Nouvelle-Aquitaine, France (MULTISTRESS project to O. Chastel), the CPER ECONAT, and INSU-EC2CO Ecodyn 2014 Program (to O. Chastel). This study has been carried out with financial support from the French National Research Agency (ANR) in the frame of the "Investments for the future" Program, within the Cluster of Excellence COTE (ANR-10-LABX-45). At the CEBC, we thank Marie Pallud and Cécile Ribout for their help in thyroids hormone assay and molecular sexing. The authors thank all the fieldworkers from Ligue Pour la Protection des Oiseaux (LPO) for their help in the long-term monitoring and ringing program (PP533) supported by the Centre de Recherches sur la Biologie des Populations d'Oiseaux (CRBPO). We also thank three anonymous reviewers for providing constructive and stimulating comments on our manuscript. This study was approved by the French Animal Ethic Committee (authorisation number: APAFIS#15629-2019032823161213).

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2020.144611>.

## References

- Barrie, L., Gregor, D., Hargrave, B.T., Lake, R., Muir, D., Shearer, R., et al., 1992. Arctic contaminants: sources, occurrence and pathways. *Sci. Total Environ.* 122, 1–74.
- Berntsen, H.F., Bjørklund, C.G., Audinot, J.-N., Hofer, T., Verhaegen, S., Lentzen, E., et al., 2017. Time-dependent effects of perfluorinated compounds on viability in cerebellar granule neurons: dependence on carbon chain length and functional group attached. *Neurotoxicol.* 63, 70–83.
- Blévin, P., Angelier, F., Tartu, S., Bustamante, P., Herzke, D., Moe, B., et al., 2017a. Perfluorinated substances and telomeres in an Arctic seabird: cross-sectional and longitudinal approaches. *Environ. Pollut.* 230, 360–367.
- Blévin, P., Tartu, S., Ellis, H.I., Chastel, O., Bustamante, P., Parenteau, C., et al., 2017b. Contaminants and energy expenditure in an Arctic seabird: organochlorine pesticides and perfluoroalkyl substances are associated with metabolic rate in a contrasted manner. *Environ. Res.* 157, 118–126.
- Blévin, P., Shaffer, S.A., Bustamante, P., Angelier, F., Picard, B., Herzke, D., et al., 2020. Contaminants, prolactin and parental care in an Arctic seabird: contrasted associations of perfluoroalkyl substances and organochlorine compounds with egg-turning behavior. *Gen. Comp. Endocrinol.* 291, 113420.
- Boisvert, G., Sonne, C., Rigét, F.F., Dietz, R., Letcher, R.J., 2019. Bioaccumulation and biomagnification of perfluoroalkyl acids and precursors in East Greenland polar bears and their ringed seal prey. *Environ. Pollut.* 252, 1335–1343.

- Braune, B.M., Trudeau, S., Jeffrey, D.A., Mallory, M.L., 2011. Biomarker responses associated with halogenated organic contaminants in northern fulmars (*Fulmarus glacialis*) breeding in the Canadian Arctic. *Environ. Pollut.* 159, 2891–2898.
- Braune, B.M., Gaston, A.J., Letcher, R.J., Grant Gilchrist, H., Mallory, M.L., Provencher, J.F., 2014. A geographical comparison of chlorinated, brominated and fluorinated compounds in seabirds breeding in the eastern Canadian Arctic. *Environ. Res.* 134, 46–56.
- Buck, R.C., Franklin, J., Berger, U., Conder, J.M., Cousins, I.T., de Voogt, P., et al., 2011. Perfluoroalkyl and polyfluoroalkyl substances in the environment: terminology, classification, and origins. *Integr. Environ. Assess. Manag.* 7, 513–541.
- Bustnes, J.O., Bakken, V., Skaare, J.U., Erikstad, K.E., 2003. Age and accumulation of persistent organochlorines: a study of arctic-breeding glaucous gulls (*Larus hyperboreus*). *Environ. Toxicol. Chem.* 22, 2173–2179.
- Bustnes, J.O., Borgå, K., Erikstad, K.E., Lorentsen, S.H., Herzke, D., 2008a. Perfluorinated, brominated, and chlorinated contaminants in a population of lesser black-backed gulls (*Larus fuscus*). *Environ. Toxicol. Chem.* 27, 1383–1392.
- Bustnes, J.O., Erikstad, K.E., Lorentsen, S.-H., Herzke, D., 2008b. Perfluorinated and chlorinated pollutants as predictors of demographic parameters in an endangered seabird. *Environ. Pollut.* 156, 417–424.
- Carravieri, A., Burthe, S.J., de la Vega, C., Yonehara, Y., Daunt, F., Newell, M.A., Jeffreys, R.M., Lawlor, A.J., Hunt, A., Shore, R.F., Pereira, M.G., Green, J.A., 2020. Interactions between environmental contaminants and gastrointestinal parasites: novel insights from an integrative approach in a marine predator. *Environ. Sci. Technol.* 54, 8938–8948.
- Cheng, J., Lv, S., Nie, S., Liu, J., Tong, S., Kang, N., et al., 2016. Chronic perfluorooctane sulfonate (PFOS) exposure induces hepatic steatosis in zebrafish. *Aquat. Toxicol.* 176, 45–52.
- Conder, J.M., Hoke, R.A., Wd, Wolf, Russell, M.H., Buck, R.C., 2008. Are PFCAs bioaccumulative? A critical review and comparison with regulatory criteria and persistent lipophilic compounds. *Environ. Sci. Technol.* 42, 995–1003.
- Corman, A.-M., Mendel, B., Voigt, C.C., Garthe, S., 2016. Varying foraging patterns in response to competition? A multicolony approach in a generalist seabird. *Ecol. Evol.* 6, 974–986.
- Costantini, D., Blevin, P., Herzke, D., Moe, B., Gabrielsen, G.W., Bustnes, J.O., et al., 2019. Higher plasma oxidative damage and lower plasma antioxidant defences in an Arctic seabird exposed to longer perfluoroalkyl acids. *Environ. Res.* 168, 278–285.
- Couderc, M., Poirier, L., Zalouk-Vergnoux, A., Kamari, A., Blanchet-Letrouve, I., Marchand, P., et al., 2015. Occurrence of POPs and other persistent organic contaminants in the European eel (*Anguilla anguilla*) from the Loire estuary, France. *Sci. Total Environ.* 505, 199–215.
- Danovaro, R., 2003. Pollution threats in the Mediterranean Sea: an overview. *Chem. Ecol.* 19, 15–32.
- Darras, V.M., Verhoelst, C.H., Reynolds, G.E., Kühn, E.R., Van der Geyten, S., 2006. Thyroid hormone deiodination in birds. *Thyroid* 16, 25–35.
- DeWitt, J., 2015. *Toxicological Effects of Perfluoroalkyl and Polyfluoroalkyl Substances*. Springer Press.
- Elliott, J.E., Elliott, K.H., 2013. Tracking marine pollution. *Science* 340, 556.
- Fernandes, A.R., Mortimer, D., Holmes, M., Rose, M., Zhihua, L., Huang, X., et al., 2018. Occurrence and spatial distribution of chemical contaminants in edible fish species collected from UK and proximate marine waters. *Environ. Int.* 114, 219–230.
- Fridolfsson, A.-K., Ellegren, H., 1999. A simple and universal method for molecular sexing of non-ratite birds. *J. Avian Biol.* 30, 116–121.
- Furness, R.W., Camphuysen, K., 1997. Seabirds as monitors of the marine environment. *ICES J. Mar. Sci.* 54, 726–737.
- Giesy, J.P., Kannan, K., 2001. Global distribution of perfluorooctane sulfonate in wildlife. *Environ. Sci. Technol.* 35, 1339–1342.
- Haarr, A., Hylland, K., Eckbo, N., Gabrielsen, G.W., Herzke, D., Bustnes, J.O., Blévin, P., Chastel, O., Moe, B., Hanssen, S.A., Sagerup, K., Borgå, K., 2018. DNA damage in Arctic seabirds: baseline, sensitivity to a genotoxic stressor, and association with organohalogen contaminants. *Environ. Toxicol. Chem.* 37, 1084–1091.
- Hanssen, S.A., Hasselquist, D., Folstad, I., Erikstad, K.E., 2005. Cost of reproduction in a long-lived bird: incubation effort reduces immune function and future reproduction. *Proc. Biol. Sci.* 272, 1039–1046.
- Herzke, D., Nygård, T., Berger, U., Huber, S., Røv, N., 2009. Perfluorinated and other persistent halogenated organic compounds in European shag (*Phalacrocorax aristotelis*) and common eider (*Somateria mollissima*) from Norway: a suburban to remote pollutant gradient. *Sci. Total Environ.* 408, 340–348.
- Jacobsen, A.V., Nordén, M., Engwall, M., Scherbag, N., 2018. Effects of perfluorooctane sulfonate on genes controlling hepatic fatty acid metabolism in livers of chicken embryos. *Environ. Sci. Pollut. Res.* 25, 23074–23081.
- Jouanneau, W., Bårdsen, B.-J., Herzke, D., Johnsen, T.V., Eulaers, I., Bustnes, J.O., 2020. Spatiotemporal analysis of perfluoroalkyl substances in white-tailed eagle (*Haliaeetus albicilla*) nestlings from Northern Norway—a ten-year study. *Environ. Sci. Technol.* 54, 5011–5020.
- Kannan, K., Tao, L., Sinclair, E., Pastva, S.D., Jude, D.J., Giesy, J.P., 2005. Perfluorinated compounds in aquatic organisms at various trophic levels in a Great Lakes food chain. *Arch. Environ. Contam. Toxicol.* 48, 559–566.
- Kelly, B.C., Ikonomou, M.G., Blair, J.D., Surridge, B., Hoover, D., Grace, R., et al., 2009. Perfluoroalkyl contaminants in an Arctic marine food web: trophic magnification and wildlife exposure. *Environ. Sci. Technol.* 43, 4037–4043.
- Land, M., de Wit, C.A., Bignert, A., Cousins, I.T., Herzke, D., Johansson, J.H., et al., 2018. What is the effect of phasing out long-chain per- and polyfluoroalkyl substances on the concentrations of perfluoroalkyl acids and their precursors in the environment? A systematic review. *Environm. Evid.* 7, 4.
- Liu, Y., Wang, J., Fang, X., Zhang, H., Dai, J., 2011. The thyroid-disrupting effects of long-term perfluorononanoate exposure on zebrafish (*Danio rerio*). *Ecotoxicology* 20, 47–55.
- Lucia, M., Strøm, H., Bustamante, P., Herzke, D., Gabrielsen, G.W., 2017. Contamination of ivory gulls (*Pagophila eburnea*) at four colonies in Svalbard in relation to their trophic behaviour. *Polar Biol.* 40, 917–929.
- Maynard, L., Davoren, G., 2020. Inter-colony and interspecific differences in the isotopic niche of two sympatric gull species in Newfoundland. *Mar. Ornithol.* 48, 103–109.
- McNabb, F.M.A., 2007. The hypothalamic-pituitary-thyroid (HPT) axis in birds and its role in bird development and reproduction. *Crit. Rev. Toxicol.* 37, 163–193.
- Melnes, M., Gabrielsen, G.W., Herzke, D., Sagerup, K., Jenssen, B.M., 2017. Dissimilar effects of organohalogenated compounds on thyroid hormones in glaucous gulls. *Environ. Res.* 158, 350–357.
- Muir, D.C.G., de Wit, C.A., 2010. Trends of legacy and new persistent organic pollutants in the circumpolar arctic: overview, conclusions, and recommendations. *Sci. Total Environ.* 408, 3044–3051.
- Muir, D.C.G., Bossi, R., Carlsson, P., Evans, M., De Silva, A., Halsall, C., et al., 2019. Levels and trends of poly- and perfluoroalkyl substances in the Arctic environment – an update. *Emerg. Contam.* 5, 240–271.
- Munoz, G., Budzinski, H., Labadie, P., 2017a. Influence of environmental factors on the fate of legacy and emerging per- and polyfluoroalkyl substances along the salinity/turbidity gradient of a macrotidal estuary. *Environ. Sci. Technol.* 51, 12347–12357.
- Munoz, G., Labadie, P., Geneste, E., Pardon, P., Tartu, S., Chastel, O., et al., 2017b. Biomonitoring of fluoroalkylated substances in Antarctica seabird plasma: development and validation of a fast and rugged method using on-line concentration liquid chromatography tandem mass spectrometry. *J. Chromatogr. A* 1513, 107–117.
- Munoz, G., Budzinski, H., Babut, M., Lobry, J., Selleslagh, J., Tapie, N., et al., 2019. Temporal variations of perfluoroalkyl substances partitioning between surface water, suspended sediment, and biota in a macrotidal estuary. *Chemosphere* 233, 319–326.
- Munsch, C., Nadege, B., Pollono, C., Aminot, Y., 2019. Perfluoroalkyl substances (PFASs) in the marine environment: spatial distribution and temporal profile shifts in shellfish from French coasts. *Chemosphere* 228, 640–648.
- Nøst, T.H., Helgason, L.B., Harju, M., Heimstad, E.S., Gabrielsen, G.W., Jenssen, B.M., 2012. Halogenated organic contaminants and their correlations with circulating thyroid hormones in developing Arctic seabirds. *Sci. Total Environ.* 414, 248–256.
- Peig, J., Green, A.J., 2009. New perspectives for estimating body condition from mass/length data: the scaled mass index as an alternative method. *Oikos* 118, 1883–1891.
- Ramhøj, L., Hass, U., Gilbert, M.E., Wood, C., Svingsen, T., Usai, D., et al., 2020. Evaluating thyroid hormone disruption: investigations of long-term neurodevelopmental effects in rats after perinatal exposure to perfluorohexane sulfonate (PFHxS). *Sci. Rep.* 10, 2672.
- Ren, X.-M., Qin, W.-P., Cao, L.-Y., Zhang, J., Yang, Y., Wan, B., et al., 2016. Binding interactions of perfluoroalkyl substances with thyroid hormone transport proteins and potential toxicological implications. *Toxicology* 366–367, 32–42.
- Sebastiano, M., Angelier, F., Blevin, P., Ribout, C., Sagerup, K., Descamps, S., et al., 2020. Exposure to PFAS is associated with telomere length dynamics and demographic responses of an arctic top predator. *Environ. Sci. Technol.* 54, 10217–10226.
- Simonnet-Laprade, C., Budzinski, H., Maciejewski, K., Le Menach, K., Santos, R., Alliot, F., et al., 2019. Biomagnification of perfluoroalkyl acids (PFAAs) in the food web of an urban river: assessment of the trophic transfer of targeted and unknown precursors and implications. *Environ Sci Process Impacts* 21, 1864–1874.
- Smits, J.E., Fernie, K.J., Bortolotti, G.R., Marchant, T.A., 2002. Thyroid hormone suppression and cell-mediated immunomodulation in American kestrels (*Falco sparverius*) exposed to PCBs. *Arch. Environ. Contam. Toxicol.* 43, 338–344.
- Sun, J., Bossi, R., Bustnes, J.O., Helander, B., Boertmann, D., Dietz, R., et al., 2019. White-tailed eagle (*Haliaeetus albicilla*) body feathers document spatiotemporal trends of Perfluoroalkyl substances in the northern environment. *Environ. Sci. Technol.* 53, 12744–12753.
- Sunderland, E.M., Hu, X.C., Dassuncao, C., Tokranov, A.K., Wagner, C.C., Allen, J.G., 2019. A review of the pathways of human exposure to poly- and perfluoroalkyl substances (PFASs) and present understanding of health effects. *J. Expos. Sci. Environ. Epidemiol.* 29, 131–147.
- Tartu, S., Gabrielsen, G.W., Blévin, P., Ellis, H., Bustnes, J.O., Herzke, D., et al., 2014. Endocrine and fitness correlates of long-chain perfluorinated carboxylates exposure in Arctic breeding black-legged kittiwakes. *Environ. Sci. Technol.* 48, 13504–13510.
- Verreault, J., Skaare, J.U., Jenssen, B.M., Gabrielsen, G.W., 2004. Effects of organochlorine contaminants on thyroid hormone levels in arctic breeding glaucous gulls, *Larus hyperboreus*. *Environ. Health Perspect.* 112, 532–537.
- Verreault, J., Houde, M., Gabrielsen, G.W., Berger, U., Haukås, M., Letcher, R.J., et al., 2005. Perfluorinated alkyl substances in plasma, liver, brain, and eggs of glaucous gulls (*Larus hyperboreus*) from the Norwegian Arctic. *Environ. Sci. Technol.* 39, 7439–7445.
- Verreault, J., Villa, R.A., Gabrielsen, G.W., Skaare, J.U., Letcher, R.J., 2006. Maternal transfer of organohalogen contaminants and metabolites to eggs of Arctic-breeding glaucous gulls. *Environ. Pollut.* 144, 1053–1060.
- Vicente, J., Sanpera, C., García-Tarrasón, M., Pérez, A., Lacorte, S., 2015. Perfluoroalkyl and polyfluoroalkyl substances in entire clutches of Audouin's gulls from the Ebro delta. *Chemosphere* 119, S62–S68.
- Wan, H.T., Zhao, Y.G., Wei, X., Hui, K.Y., Giesy, J.P., Wong, C.K.C., 2012. PFOS-induced hepatic steatosis, the mechanistic actions on  $\beta$ -oxidation and lipid transport. *Biochim. Biophys. Acta Gen. Subj.* 1820, 1092–1101.
- Wang, Z., DeWitt, J.C., Higgins, C.P., Cousins, I.T., 2017. A never-ending story of per- and polyfluoroalkyl substances (PFASs)? *Environ. Sci. Technol.* 51, 2508–2518.
- Weiss, J.M., Andersson, P.L., Lamoree, M.H., Leonard, P.E.G., van Leeuwen, S.P.J., Hamers, T., 2009. Competitive binding of poly- and perfluorinated compounds to the thyroid hormone transport protein transthyretin. *Toxicol. Sci.* 109, 206–216.

- Wong, F., Shoeib, M., Katsoyiannis, A., Eckhardt, S., Stohl, A., Bohlin-Nizzetto, P., et al., 2018. Assessing temporal trends and source regions of per- and polyfluoroalkyl substances (PFASs) in air under the Arctic Monitoring and Assessment Programme (AMAP). *Atmos. Environ.* 172, 65–73.
- Xie, S., Wang, T., Liu, S., Jones, K.C., Sweetman, A.J., Lu, Y., 2013. Industrial source identification and emission estimation of perfluorooctane sulfonate in China. *Environ. Int.* 52, 1–8.
- Xu, J., Shimpi, P., Armstrong, L., Salter, D., Slitt, A.L., 2016. PFOS induces adipogenesis and glucose uptake in association with activation of Nrf2 signaling pathway. *Toxicol. Appl. Pharmacol.* 290, 21–30.
- Yeung, L.W.Y., Guruge, K.S., Yamanaka, N., Miyazaki, S., Lam, P.K.S., 2007. Differential expression of chicken hepatic genes responsive to PFOA and PFOS. *Toxicology* 237, 111–125.