

Lithium isotopes in marine food webs

FANNY THIBON¹, LUCAS WEPPE¹, PACO BUSTAMANTE², FRANÇOIS OBERHÄNSLI³, MARC METIAN³, CARINE CHURLAUD², MARYLINE MONTANES¹, THOMAS LACOUÉ-LABARTHE², PHILIPPE TELOUK⁴, YVES CHEREL⁵ AND NATHALIE VIGIER⁶

¹LOV- CNRS-Sorbonne Université

²LIENSs - CNRS-La Rochelle Université

³IAEA-REL

⁴ENS Lyon

⁵CEBC - CNRS-La Rochelle Université

⁶LOV - CNRS - Sorbonne Université

Presenting Author: thibon.fanny@orange.fr

Lithium (Li) production has massively increased over the past decades and is essentially used for batteries, ceramics, and medicine [1-2]. However, Li concentrations and its isotopic compositions ($\delta^7\text{Li}$) are poorly documented in aquatic ecosystems, despite its accumulation in marine organisms, especially at the base of the trophic webs [3]. Here, we present field and laboratory studies reporting $\delta^7\text{Li}$ in various organs and whole organisms of marine plankton, bivalves, cephalopods, crustaceans, and fish collected in different biogeographical regions (the Mediterranean Sea, Bay of Biscay (BB), New Caledonia (NC), and Kerguelen Islands). The main aim of the study was to assess the role of ecological and environmental parameters on $\delta^7\text{Li}$ values.

$\delta^7\text{Li}$ values ranged from 4.6 ‰ (digestive gland of bivalves) to 31.7 ‰ (>35 μm plankton). Seawater $\delta^7\text{Li}$ being homogeneously distributed at ~ 31.2 ‰, marine organisms thus fractionate Li isotopes in favor of the light isotope (^6Li). There are no trends in $\delta^7\text{Li}$ values with the trophic position. However, when the same organs from similar organisms (bivalves, cephalopods, crustaceans, and fish) are compared, lower $\delta^7\text{Li}$ are systematically found in NC compared to BB, overall suggesting an influence of the environmental conditions on $\delta^7\text{Li}$. Moreover, cultured organisms Li isotope fractionation increases when facing high Li levels in water.

Within the same taxonomic group, significant differences were also observed among organs with the lowest $\delta^7\text{Li}$ values in gills for fish and in the digestive gland for bivalves. This indicates a key role of physiology on Li isotopes distribution. A trend with Na contents supports an intimate link between Li transport and regulation and the activity of Na-H Exchangers [4]. Overall, the work highlights the interest of $\delta^7\text{Li}$ to investigate Li ecotoxicology and physiology. It may also help to unravel the magnitude of biological activity (“vital effects”) on $\delta^7\text{Li}$ measured in fossil biogenic carbonates.

[1] Choi et al. (1998), *Nat. Commun.* **10**, 5371–5378. [2] L  gu  rinel et al. (2018), *COMES seminar*. [3] Thibon and Weppe et al. (2020), *Stoten*, **751**, 141453. [4] Counillon et al. (2016), *Biochim. Biophys. Acta.* **1863**, 2465–2480