

## Food supplementation of the Montpellier snake *Malpolon monspessulanus* in the wild

GREGORY DESO<sup>1\*</sup>, ALOYS CROUZET<sup>1</sup> & XAVIER BONNET<sup>2</sup>


<sup>1</sup>AHPAM - Association Herpétologique de Provence Alpes Méditerranée, Maison des Associations, 384 route de Caderousse, 84100 Orange, France

<sup>2</sup>Centre d'Étude Biologique de Chizé, UMR-7372, CNRS-Université de La Rochelle, 79360, Villiers en Bois, France

\*Corresponding author e-mail: ahpam.contact@gmail.com

Under natural settings, a variety of snake species are occasional scavengers, especially piscivorous snakes and pit vipers (DeVault & Krochmal, 2002). In the case of the Montpellier snake *Malpolon monspessulanus* the species has been observed both eating bait intended for other animals and roadkill (Valverde, 1974; Ventura, 2012). Several researchers have taken advantage of the propensity of wild snakes to accept carrion (e.g. mammals, fish) offered by humans to address a variety of ecological questions (Sazima & Strüßmann, 1990; Marques & Sazima, 1997; Wasko & Sasa, 2012; Glaudas & Alexander, 2017). Further investigation of the foraging response of wild snakes to carrion offered by researchers is of interest because it has the potential to improve animal welfare, for example to place electronic devices without the need for force-feeding (Shine, 1987). It could also be used to supplement the food intake of wild snakes, thereby promoting reproduction and recruitment in fragile populations (Taylor et al., 2005). Weitzmann & Pretus (2018) made an unsuccessful (prey not consumed despite snake interest) attempt at food supplementation with the Montpellier snake and this led us to undertake a feeding test in 2021 in Bédarrides, Vaucluse district, France, with two large free-ranging male Montpellier snakes both in the range of 1.7-1.8 m long and 1.5 to 1.6 kg body mass.

We offered different food items (approximately 20-110 g). The mass of the meals was estimated to be in the range of



**Figure 1.** Setting food supplementation tests for Montpellier snakes - **A.** A camera trap (Num'axes) was positioned by food deposited in a place frequently used by snakes, **B.** A successful test, the snake swallowed the chicken leg (BHS video, 2022)

the prey commonly consumed by large Montpellier snakes (e.g., small to large rodents). We placed the food items in locations where snakes were frequently observed during the study and used a video trap to monitor the snakes' behaviour (Num'Axes Trail Camera PIE 1023) (Fig. 1). We tested three types of easily available food items: chicken legs, quail eggs, and chipolata (pork) sausages (Table 1). Chicken legs were presented alone, eggs and sausage together. Tests where the bait was quickly covered by ants were discarded.

Chicken legs were tested five times with the first snake and once with the second snake (Table 1). The first snake found

**Table 1.** Details of dietary supplementation tests with Montpellier snakes (#1 and #2) using different food items

Snake #	Food type	Date	Time of various activities				Test outcome
			Food deposited	Snake detects food	Swallowing		
					Begins	Ends	
1	Chicken leg	18 Jun	11:44	12: 22	13:15	13: 36	Successful
1	Chicken leg	29 Aug	10:44	-	13:42	14: 00	Successful
1	Chicken leg	5 Sept	14:00	-	17:00	NA	Successful
1	Chicken leg	16 Jul	15:00	-	-	-	Unsuccessful
1	Chicken leg	13 Sept	13:10	-	-	-	Unsuccessful
1	Quail egg + chipolata sausage	30 Jun	17:00	-	-	-	Unsuccessful
2	Chicken leg	1 Sept	11:35	14:10	-	-	Unsuccessful
2	Quail egg + chipolata sausage	24 Jun	10:10	11: 05	-	-	Unsuccessful

and ate chicken legs on three occasions (BHS video, 2022) but rejected it twice while the second snake twice showed interest in the chicken leg but did not eat it on either occasion (Table 1). Both snakes were tested once with the egg and sausage combination but neither consumed the food.

Video recording is an essential means for describing behaviours that are difficult to observe and quantify in the field (Ballouard et al., 2016; Clark, 2006; Glaudas et al., 2017) and on this occasion has been used to reveal the first successful food supplementation for a wild Montpellier snake. We recommend using video monitoring to assess potential benefits and disadvantages of food supplementation. This process, which allows remote monitoring from a mobile phone, is a technology that is not yet widely used for behavioural monitoring of snakes. However, it is particularly appropriate because i) most wild snakes are scared by humans and are reluctant to eat in their presence and ii) it is essential to determine which animal has eaten the food supplement (e.g., the targeted snake species, scavengers, or potential snake predators).

## ACKNOWLEDGEMENTS

We thank Xavier Glaudas, Nicolas Dubos, and Ivan Sazima for useful comments.

## REFERENCES

- Ballouard, J.M., Bonnet, X., Gravier, C., Ausanneau, M. & Caron, S. (2016). Artificial water ponds and camera trapping of tortoises, and other vertebrates, in a dry Mediterranean landscape. *Wildlife Research* 43: 533-543. doi:10.1071/WR16035
- BHS video (2022). Food supplementation of a Montpellier snake (*Malpolon monspessulanus*) with a chicken leg. <https://youtu.be/xKtBPMooFa0>
- Clark, R. (2006). Fixed videography to study predation behavior of an ambush foraging snake, *Crotalus horridus*. *Copeia* 2006(2): 181–187.
- DeVault, T.L. & Krochmal, A.R. (2002). Scavenging by snakes: an examination of the literature. *Herpetologica* 58(4): 429–436.
- Glaudas, X. & Alexander, G.J. (2017). Food supplementation affects the foraging ecology of a low-energy, ambush-foraging snake. *Behavioral Ecology and Sociobiology* 71: 5.
- Marques, O.A.V. & Sazima, I. (1997). Diet and feeding behavior of the Coral Snake, *Micrurus corallinus*, from the Atlantic forest of Brazil. *Herpetological Natural History* 5: 88–93.
- Sazima, C. & Strüssmann, I. (1990). Necrofagia em serpentes brasileira: Exemplos e previsões. Re-vista *Brasileira de Biologia* 50: 463–468.
- Shine, R. (1987). Intraspecific variation in thermoregulation, movements and habitat use by Australian blacksnakes, *Pseudechis porphyriacus* (Elapidae). *Journal of Herpetology* 21: 165–177.
- Taylor, E.N., Malawy, M.A., Browning, D.M., Lemar, S.V. & DeNardo, D.F. (2005). Effects of food supplementation on the physiological ecology of female western diamond-backed rattlesnakes (*Crotalus atrox*). *Oecologia* 144(2): 206–213.
- Valverde, J.A. (1974). *Malpolon monspessulanus* llevando *Lacerta lepida* aplasta dopor un coche. *Doñana, Acta Vertebrata* 1: 56.
- Ventura, F. (2012). Comportamiento carroñero en *Malpolon monspessulanus*. *Boletín de la Asociación Herpetológica Española*, 23(1): 8–10 pp.
- Wasko, D.K. & Sasa, M. (2012). Food resources influence spatial ecology, habitat selection, and foraging behavior in an ambush-hunting snake (Viperidae: *Bothrops asper*): an experimental study. *Zoology* 115: 179–187.
- Weitzmann, B. & Pretus, F. (2018). Cita de comportament carronyaire de *Malpolon monspessulanus* obtinguda mitjançant trampeig fotogràfic a Catalunya. *Bulletí de la Societat Catalana d'Herpetologia* 25: 7–10.

Accepted: 2 December 2021